



Bottomonium and exotic spectroscopy

22nd Flavor Physics and CP Violation (FPCP)

Renu


On the behalf of Belle II Collaboration

Supported by US DOE funding


27th May, 2024 - 31st May, 2024




Bottomonium Spectrum



- ▶ Below the $B\bar{B}$ threshold states are well described by potential models.
- ▶ Above $B\bar{B}$ threshold states exhibit unexpected properties:
 - ◆ Hadronic transitions to lower bottomonia are strongly enhanced.
 - ◆ The η transitions are not suppressed compared to $\pi^+\pi^-$ transitions. Strong violation of Heavy Quark Spin Symmetry.
 - ◆ $Z_b^+(10610)$ or $Z_b^+(10650)$: observed near the $B^{(*)}\bar{B}^*$ thresholds, properties are consistent with $B^{(*)}\bar{B}^*$ molecules.




Exotic: molecule, compact tetra-quark.



- ▶ Conventional bottomonium (pure $b\bar{b}$ state)
- ▶ Bottomonium like states (mix of $b\bar{b}$ and $B\bar{B}$)
- ▶ Purely exotic states (Z_b)

Belle (II) relevant datasets





Bottomonium below $B\bar{B}$ threshold




Search for $h_b(2P) \rightarrow \Upsilon(1S)\eta$ and $h_b(1P,2P) \rightarrow \Upsilon(1S)\pi^0$ at Belle



- ▶ The properties of spin-singlet $h_b(1P,2P)$ are expected to be similar to spin-triplet partners $\chi_{b1}(1P,2P)$ state.
- ▶ **Theoretical prediction:** the ratio of the annihilation rates for the $h_b(1P)$ and $h_b(2P)$ is the same as the corresponding ratio for $\chi_{b1}(1P)$ and $\chi_{b1}(2P)$,
 $R_{h_b} = R_{\chi_{b1}}$. PRD 86, 094013 (2012)
- ▶ Based on current results, the $R_{h_b}/R_{\chi_{b1}} = 0.24^{+0.47}_{-0.24}$ with 1.5σ discrepancy from unity. This discrepancy will increase if the rate of $h_b(2P) \rightarrow \Upsilon(1S)\eta$ is as large as 10%



Preliminary results!





- ▶ Evidence for $h_b(2P) \rightarrow \Upsilon(1S)\eta$ with 3.5σ significance.
- ▶ $\mathcal{B}(h_b \rightarrow \Upsilon(1S)\eta) = (7.1_{-3.2}^{+3.5} \pm 0.8) \times 10^{-3}$
- ▶ No significant $h_b(1P, 2P) \rightarrow \Upsilon(1S)\pi^0$ signal is observed.
- ▶ Upper limits at the 90% C.L. are set.
- ▶ $\mathcal{B}(h_b(1P,2P) \rightarrow \Upsilon(1S)\pi^0) < 1.8 \times 10^{-3}$ at 90% C.L

Search for $h_b(2P) \rightarrow \gamma\chi_{bJ}(1P)$ at Belle



- ▶ $h_b(2P) \rightarrow \gamma\chi_{bJ}(1P)$ is highly suppressed due to heavy quark spin flip.
- ▶ Relativized quark model predicts, $\mathcal{B}(h_b(2P) \rightarrow \gamma\chi_{bJ}(1P)) = 10^{-6} - 10^{-5}$
- ▶ According to coupled channel effect, $\mathcal{B}(h_b(2P) \rightarrow \gamma\chi_{bJ}(1P)) = 10^{-2} - 10^{-1}$ [PRD 32, 189 \(1985\)](#)
- ▶ **Experimental results needed !!** [PLB 760, 417 \(2016\)](#)



Search for $h_b(2P) \rightarrow \gamma\chi_{bJ}(1P)$ at Belle




Preliminary results!

- ▶ No significant $h_b(2P) \rightarrow \gamma\chi_{bJ}(1P)$ signal is observed.
- ▶ Upper limits at the 90% C.L. are set.

TABLE IV. Observed upper limits at 90% CL for the branching fractions of the investigated transitions.

Channel	\mathcal{B}
$h_b(2P) \rightarrow \gamma\chi_{b2}(1P)$	$< 1.2 \times 10^{-2}$
$h_b(2P) \rightarrow \gamma\chi_{b1}(1P)$	$< 5.4 \times 10^{-3}$
$h_b(2P) \rightarrow \gamma\chi_{b0}(1P)$	$< 2.7 \times 10^{-1}$



Results are consistent with the Relativized Quark Model (RQM)

Hidden flavor cross section


Discovery of $\Upsilon(10753)$

► $\Upsilon(10753)$ was observed in energy dependence of $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1,2,3$) cross sections by Belle.


[JHEP 10 \(2019\) 220](#)

► The global significance is 5.2σ

	$\Upsilon(5S)$	$\Upsilon(6S)$	New structure
M (MeV/c ²)	$10885.3 \pm 1.5^{+2.2}_{-0.9}$	$11000.0^{+4.0}_{-4.5} \ ^{+1.0}_{-1.3}$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
Γ (MeV)	$36.6^{+4.5}_{-3.9} \ ^{+0.5}_{-1.1}$	$23.8^{+8.0}_{-6.8} \ ^{+0.7}_{-1.8}$	$35.5^{+17.6}_{-11.3} \ ^{+3.9}_{-3.3}$



► $e^+e^- \rightarrow b\bar{b}$ cross section in bottomonium energy region based on the Belle and BABAR measurement.




◆ A dip near 10.75 GeV likely caused by interference between BW and smooth component.

[CPC 44, 8, 083001 \(2020\)](#)

Fit function: 3 BW+smooth component

$\Upsilon(10753)$: theoretical interpretation



Possible interpretations:

► Conventional bottomonium?

Phys. Rev. D 105, 114041 (2022)
Phys. Rev. D 106, 094013 (2022)
Phys. Rev. D 105, 074007 (2022)

► Hybrid state?

Phys. Rept. 873, 1 (2020)
Phys. Rev. D 104, 034019 (2021)

► Tetraquark state?

Phys. Rev. D 103, 074507 (2021)
Phys. Rev. D 107, 094515 (2023)



► Hadronic molecule with a small admixture of a bottomonium?

► Mass does not match $\Upsilon(3D)$ theoretical predictions, and D -wave states are not seen in e^+e^- collisions.

► $\Upsilon(4S) - \Upsilon(3D)$ mixing can be enhanced due to hadronic loops.

Unique data with energy scan near $\sqrt{s} = 10.75$ GeV

- ▶ Belle II / SuperKEKB performed an energy scan in November 2021 with a total luminosity of 19 fb^{-1} .
- ▶ **Physics Goals:**
 - ▶ Confirm and study the $\Upsilon(10753)$.
 - ▶ Improve the precision of exclusive cross-section below the $\Upsilon(5S)$.



- ▶ Belle II collected data in the gaps between the Belle points.
- ▶ The point with the highest statistics (9.8 fb^{-1}) is near the $\Upsilon(10753)$ peak.




Search for $\Upsilon(10753) \rightarrow \pi^+\pi^-\Upsilon(nS)$ at Belle II



- Discovery mode of the $\Upsilon(10753)$ (Next few slides will cover):
 - ◆ Confirm its existence
 - ◆ Measure the di-pion spectrum
 - ◆ Look for $Z_b^+(10610)$ or $Z_b^+(10650)$ intermediate contributions

Confirm $\Upsilon(10753)$ existence

- Clear signal for $\Upsilon(1S)\pi^+\pi^-$ and $\Upsilon(2S)\pi^+\pi^-$ decay mode.
- No evidence of $\Upsilon(3S)\pi^+\pi^-$




Confirm $\Upsilon(10753)$ existence

- New measurement **confirms previous Belle result**: cross section is peaking near 10.75 GeV.

	Belle + Belle II (MeV)	Belle (MeV)
$M_{\Upsilon(10753)}$	$10756.6 \pm 2.7 \pm 0.9$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
$\Gamma_{\Upsilon(10753)}$	$29.0 \pm 8.8 \pm 1.2$	$35.5^{+17.6+3.9}_{-11.3-3.3}$

- Results are consistent with the Belle results.
- Uncertainties are improved by a factor of two from previous Belle results.





Resonant structure in $\Upsilon(10753) \rightarrow \pi^+\pi^-\Upsilon(nS)$

$Z_b^+(10610)$ or $Z_b^+(10650)$ intermediate resonances



► No signal of intermediate $Z_b^+(10610)$ or $Z_b^+(10650)$ resonances are observed.

$$\Delta M_\pi = M(\pi^\pm\mu^+\mu^-) - M(\mu^+\mu^-)$$



Di-pion spectrum

- $\pi^+\pi^-\Upsilon(1S)$: $M(\pi^+\pi^-)$ distribution is consistent with phase space.
- $\pi^+\pi^-\Upsilon(2S)$: larger values of $M(\pi^+\pi^-)$ enhanced (similar to $\Upsilon(2S) \rightarrow \pi^+\pi^-\Upsilon(1S)$ process)



► Theory:

◆ Mixed $4S - 3D$ model suggests $\Upsilon(10753) \rightarrow \omega \chi_{bJ}(1P)$ could be enhanced. [PRD 104, 034036 \(2021\)](#)

► Charmonium sector:

◆ Similar to $\Upsilon(10753)$ in $e^+e^- \rightarrow \pi^+\pi^-\Upsilon(nS)$, $Y(4260)$ was observed in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ cross section by BESIII.

● Expect similar nature of $\Upsilon(10753)$ and $Y(4260)$.



◆ $Y(4260)$ was also observed in $\omega \chi_{c0}(1P)$ and $\gamma X(3872)$ by BESIII.

◆ Inspired by decay modes of $Y(4260)$ charmonium state, we expect



● $\Upsilon(10753) \rightarrow \omega \chi_{bJ}(1P)$

● $\Upsilon(10753) \rightarrow \gamma X_b$

X_b : bottomonium analogue of $X(3872)$



Search in $e^+e^- \rightarrow (\pi^+\pi^-\pi^0) \gamma \Upsilon(1S)$ process



Observation of $\Upsilon(10753) \rightarrow \omega \chi_{bJ}(1P)$ at Belle II




PRL 130, 091902 (2023)

The $e^+e^- \rightarrow \omega \chi_{bJ}(1P)$ ($J = 1,2$) cross sections peak at $\Upsilon(10753)$.

$$\Rightarrow \frac{\sigma(e^+e^- \rightarrow \omega \chi_{bJ})}{\sigma(e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-)} \sim \begin{cases} 1.5 \text{ at } \Upsilon(10753) \text{ GeV} \\ 0.15 \text{ at } \Upsilon(5S) \text{ GeV} \end{cases}$$

$\Rightarrow \Upsilon(10753)$ and $\Upsilon(5S)$ have different internal structure?



Solution 1: constructive interference

Solution II: destructive interference

Measured ratio:

$$\frac{\sigma(\Upsilon(10753) \rightarrow \omega \chi_{b1})}{\sigma(\Upsilon(10753) \rightarrow \omega \chi_{b2})} = 1.3 \pm 0.6$$

Prediction for a pure D -wave state: 15 [PLB 738, 172 \(2014\)](#)

Prediction for a $4S - 3D$ mixed state: 0.18 - 0.22

[PRD 104, 034036 \(2021\)](#)

Channel	\sqrt{s} (GeV)	N^{sig}	$\sigma_{\text{Born}}^{(\text{UL})}$ (pb)
$\omega \chi_{b1}$	10.745	$68.9^{+13.7}_{-13.5}$	$3.6^{+0.7}_{-0.7} \pm 0.4$
$\omega \chi_{b2}$		$27.6^{+11.6}_{-10.0}$	$2.8^{+1.2}_{-1.0} \pm 0.5$
$\omega \chi_{b1}$	10.805	$15.0^{+6.8}_{-6.2}$	1.6 @90% C.L.
$\omega \chi_{b2}$		$3.3^{+5.3}_{-3.8}$	1.5 @90% C.L.

Disagreement with both pure D wave state

Tension with the $4S - 3D$ mixed model (1.8σ)

Search for $\Upsilon(10753) \rightarrow \gamma X_b$ at Belle II




PRL 130, 091902 (2023)

The X_b is posited bottomonium counterpart of X(3872).

- ▶ No significant signal of X_b signal is observed.
- ▶ Upper limits on cross sections are set for $M(X_b) \in (10.45 - 10.65)$ GeV

\sqrt{s} GeV	$\sigma_B(e^+e^- \rightarrow \gamma X_b) \times \mathcal{B}(X_b \rightarrow \omega\Upsilon(1S))$
10.653	(0.14-0.55) pb
10.701	(0.25-0.84) pb
10.745	(0.06-0.14) pb
10.805	(0.08-0.37) pb



Reflection of
 $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$

► Motivation:



- Theoretically, tetra-quark interpretation predicts, a strong enhancement of the decay $\omega \eta_b(1S)$ compared to $\pi^+ \pi^- \Upsilon(nS)$ [CPC 43 \(2019\) 12, 123102](#)
- $4S - 3D$ mixed model predicts that decay rate of $\omega \eta_b(1S)$ is smaller than $\pi^+ \pi^- \Upsilon(nS)$ by a factor of 0.2-0.4 [PRD 109, 014039 \(2024\)](#)

► Strategy

◆ Partial reconstruction:

- Reconstructed ω meson in $\pi^+ \pi^- \pi^0$ and use the recoil mass of ω as signal variable

$$M_{\text{recoil}}(\pi^+ \pi^- \pi^0) = \sqrt{\left(\frac{\sqrt{s} - E^*}{c^2}\right)^2 - \left(\frac{p^*}{c}\right)^2}$$



Search for $\Upsilon(10753) \rightarrow \omega\eta_b(1S)$ at Belle II



PRD 109, 072013 (2024)

- ▶ No significant $\omega\eta_b(1S)$ signal is observed.
- ▶ Upper limits at the 90% C.L. on the Born cross section are set.
- ▶ $\sigma(e^+e^- \rightarrow \omega\eta_b(1S)) < 2.5$ pb

Ratio:

$$\frac{\sigma(\omega\eta_b)}{\sigma(\pi^+\pi^-\Upsilon(nS))} < 1.25$$

▶ Prediction for a tetra quark model: ~ 30 [CPC 43 \(2019\) 12, 123102](#)


▶ Prediction for a $4S - 3D$ mixed state: 0.2 - 0.4

[PRD 109, 014039 \(2024\)](#)

Evidence against the tetraquark model predictions.

Compatible with $S - D$ mixed model

$\omega \rightarrow \pi^+\pi^-\pi^0$ recoil mass distributions





Open flavor cross-section

Belle results


Motivation:

- The open flavor final states ($B^{(*)}\bar{B}^{(*)}$) make dominant contribution to $b\bar{b}$ cross-section.
 - Their measurements are critical for understanding the structure of $b\bar{b}$ states.
- The measured cross sections can be used in the coupled channel analysis of all available scan data to extract the parameters of the Υ states.
- Belle measured the energy dependencies of $\sigma(e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)})$ and observed an oscillatory behavior.**
 - Channels $B^{(*)}\bar{B}^{(*)}$ saturate the cross-section below the $B_s^*\bar{B}_s^*$ threshold.
- To improve the accuracy below $\Upsilon(5S)$ and understand the nature of $\Upsilon(10753)$, need more data: Belle II



Preliminary results!

- ▶ The obtained cross sections at four energies are consistent with the Belle results.
- ▶ $\sigma(e^+e^- \rightarrow B^*\bar{B}^*)$ increases rapidly above $B^*\bar{B}^*$ threshold
- ◆ Similar phenomenon was observed near $D^*\bar{D}^*$ threshold.
- ◆ **Possible interpretation:** resonance or bound state ($B^*\bar{B}^*$ or $b\bar{b}$) near $B^*\bar{B}^*$ threshold
- ◆ Inelastic channels [$\pi^+\pi^-\Upsilon(nS)$ and $\eta h_b(1P)$] could also be enhanced. Need more data to study these transitions.





Solid curve – combined Belle + Belle II data fit

Dashed curve – Belle data fit only

Preliminary results!

Comparison of $\sigma_{b\bar{b}}$ and $\sigma_{B\bar{B}} + \sigma_{B\bar{B}^*} + \sigma_{B^*\bar{B}^*}$



Black dots: Belle + BaBar
[PRL 102, 012001 (2009),
PRD 93, 011101 (2016),
CPC 44, 083001 (2020)]

Open blue circles: Belle
[JHEP 06, 137 (2021)]

Filled red circles: Belle II
[this work]




► Saturate the $\sigma_{b\bar{b}}$ cross-section below the $B_s^{(*)}\bar{B}_s^{(*)}$ threshold.

► Previously observed deviation at high energy is presumably due to $B_s^{(*)}\bar{B}_s^{(*)}$, multi-body $B^{(*)}\bar{B}^{(*)}\pi(\pi)$, etc.

Production of prompt J/ψ and Υ mesons

Production of prompt J/ψ and Υ mesons

Studied the production of prompt $J/\psi - \Upsilon$ mesons in pp collisions at $\sqrt{s} = 13$ TeV




	$J/\psi - \Upsilon(1S)$	$J/\psi - \Upsilon(2S)$
σ	133 ± 22 (stat) ± 7 (syst) ± 3 (\mathcal{B})	76 ± 21 (stat) ± 4 (syst) ± 7 (\mathcal{B})
σ_{eff}	26 ± 5 (stat) ± 2 (syst) $_{-3}^{+22}$ (theo)	14 ± 5 (stat) ± 1 (syst) $_{-1}^{+7}$ (theo)

Effective cross-sections are compatible with measurements using other particle productions.

Summary

- ▶ The understanding of the physics of highly excited heavy bottomonium is very incomplete.
- ▶ First energy scan results from Belle II are quite interesting.
- ▶ No clear indication on the nature of $\Upsilon(10753)$.
 - ◆ Improved results for mass and width of $\Upsilon(10753)$ using $\Upsilon(10753) \rightarrow \Upsilon(nS)\pi^+\pi^-$.
 - ◆ $S - D$ model compatible with $\Upsilon(10753) \rightarrow \omega\eta_b(1S)$ but not with $\Upsilon(10753) \rightarrow \omega\chi_{b1,2}(1P)$.
 - ◆ No signal of intermediate $Z_b^+(10610)$ or $Z_b^+(10650)$ resonances are observed.
- ▶ Effective cross-sections of $J/\psi - \Upsilon$ production are consistent with other particle productions measurements.

Present status






Introduction

Quark model:

M. Gell-Mann, Phys.Lett. 8, 214 (1964)

Classification scheme for hadrons in terms of valance quarks.


Hadrons are composed of mesons ($q\bar{q}$, $qq\bar{q}\bar{q}$, ...) and baryons (qqq , $qqqq\bar{q}$, ...).



- ▶ $q\bar{q}$ spectroscopy with heavy quark (mostly c or b) are best place to study quark model.
- ▶ Simple two body system, non-relativistic and narrow (with OZI suppression).
- ▶ Further, one can search for exotics with them.

Pentaquark:

S=+1 Baryon




Glueball

Color-singlet multi-gluon bound state




H-diBaryon

tightly bound 6-quark state




Tetraquark mesons

tightly bound diquark-diantiquark



loosely bound meson-antimeson "molecule"



$q\bar{q}$ -gluon hybrid mesons



Not observed in conventional matter. However, they should be allowed.

Belle II detector


- ▶ Asymmetric e^+e^- collider
- ▶ **Collected data**
 - $\sim 362 \text{ fb}^{-1}$ at Y(4S)
 - 42 fb^{-1} off-resonance, 60 MeV below Y(4S).
 - 19 fb^{-1} energy scan between 10.6 to 10.8 GeV for exotic hadron studies.

Features:

- ▶ Near-hermetic detector
- ▶ Excellent vertexing and tracking
- ▶ High-efficiency detection of neutrals (γ , π^0 , η , η' , ...)
- ▶ Good charged particle reconstruction.


Record-breaking instantaneous luminosity:

$$4.7 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$$



Coupled channel analysis

Hüsken, Mitchell, Swanson, PRD 106, 094013 (2022)



All available scan data

K-matrix: scattering via $\Upsilon(4S)$, $\Upsilon(10753)$, $\Upsilon(5S)$, $\Upsilon(6S)$ or non-resonantly.

Results: pole positions, branching fraction, energy dependence of scattering amplitudes.

Accuracy above $\Upsilon(6S)$ and near $\Upsilon(10753)$ is poor.

Energy dependence of $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ cross section

Decay modes used:

$B^+ \rightarrow$	$B^0 \rightarrow$
$\bar{D}^0 \pi^+$	$D^- \pi^+$
$\bar{D}^0 \pi^+ \pi^+ \pi^-$	$D^- \pi^+ \pi^+ \pi^-$
$\bar{D}^{*0} \pi^+$	$D^{*-} \pi^+$
$\bar{D}^{*0} \pi^+ \pi^+ \pi^-$	$D^{*-} \pi^+ \pi^+ \pi^-$
$D_s^+ \bar{D}^0$	$D_s^+ D^-$
$D_s^{*+} \bar{D}^0$	$D_s^{*+} D^-$
$D_s^+ \bar{D}^{*0}$	$D_s^+ D^{*-}$
$D_s^{*+} \bar{D}^{*0}$	$D_s^{*+} D^{*-}$
$J/\psi K^+$	$J/\psi K_S$
$J/\psi K_S \pi^+$	$J/\psi K^+ \pi^-$
$J/\psi K^+ \pi^+ \pi^-$	
$D^- \pi^+ \pi^+$	$D^{*-} K^+ K^- \pi^+$
$D^{*-} \pi^+ \pi^+$	


$D^0 \rightarrow$	$D^+ \rightarrow$	$D_s^+ \rightarrow$
$K^- \pi^+$	$K^- \pi^+ \pi^+$	$K^+ K^- \pi^+$
$K^- \pi^+ \pi^0$	$K^- \pi^+ \pi^+ \pi^0$	$K^+ K_S$
$K^- \pi^+ \pi^+ \pi^-$	$K_S \pi^+$	$K^+ K^- \pi^+ \pi^0$
$K_S \pi^+ \pi^-$	$K_S \pi^+ \pi^0$	$K^+ K_S \pi^+ \pi^-$
$K_S \pi^+ \pi^- \pi^0$	$K_S \pi^+ \pi^+ \pi^-$	$K^- K_S \pi^+ \pi^+$
$K^+ K^-$	$K^+ K^- \pi^+$	$K^+ K^- \pi^+ \pi^+ \pi^-$
$K^+ K^- K_S$		$K^+ \pi^+ \pi^-$
		$\pi^+ \pi^+ \pi^-$

Energy dependence of $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ cross section

► Method:

- ◆ Reconstruct one B in full hadronic channels.
- ◆ Key variables for analysis are
 - ◆ $M_{bc} = \sqrt{(E_{cm}/2)^2 - p_B^2}$
 - ◆ $\Delta E' = \Delta E - M_{bc} + M_B$, where $\Delta E = E_B - E_{cm}/2$
- ◆ $\Delta E'$ has improved resolution and allows all desired two-body decays to be selected with a common cut
- ◆ Populations of each can be studied by fitting the projections onto the M_{bc} axis for all energies at which data were accumulated
- ◆ $B^* \rightarrow B\gamma$ decays are not reconstructed.





$\Delta E'$ vs M_{bc} at $E_{cm} = 10.746$ GeV



Energy dependence of $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ cross section

M_{bc} fit at scan energies

- ▶ M_{bc} fit distribution:
- ▶ $\Delta E'$ signal region (upper)
- ▶ $\Delta E'$ side-bands (lower)
- ▶ $e^+e^- \rightarrow B\bar{B}, B\bar{B}^*, B^*\bar{B}^*$ signals at $\sqrt{s} \sim 10.75$ GeV can be clearly observed
- ▶ Contribution of $\Upsilon(4S) \rightarrow B\bar{B}$ production via ISR is visible well (black dotted histograms)
- ▶ At $\sqrt{s} = 10.653$ GeV, the sharp cut of the data at right edge is due to threshold effect



Bottomonium (-like) at Belle II

► Four ways to access bottomonia:

◆ **Direct production** from e^+e^- : $J^{PC} = 1^{--}$: $\Upsilon(nS)$


◆ **ISR production**: $J^{PC} = 1^{--}$: $\Upsilon(nS)$

◆ **Hadronic transitions** from $\Upsilon(nS)$ through $\eta, \pi\pi, \dots$


$$J^{PC} = 0^{-+}, 1^{--}, 1^{+-} \dots : \Upsilon(nS), \eta_b(nS), h_b(nS), \dots$$

◆ **Radiative transitions** from $\Upsilon(nS)$

$$J^{PC} = 0^{-+}, 0^{++}, 1^{++}, 2^{++} : \eta_b(nS), \chi_b(nP)$$



ISR production



Hadronic transitions

