


# Searches for invisible new particles at Belle II

21<sup>st</sup> Lomonosov Conference on Elementary Particle Physics

Moscow State University  
August 24-30, 2023

**Luigi Corona** - INFN Pisa  
on behalf of the Belle II collaboration  
 [luigi.corona@pi.infn.it](mailto:luigi.corona@pi.infn.it)





# Dark matter searches

## Dark Matter (DM)

- It is one of the most compelling phenomena in support for physics beyond the Standard Model (SM)
  - It exists, awaiting for discovery


Albada et al., *Astrophysical Journal* (1985)




## How to search for DM?

- Focus on searches at **collider** experiment (Belle II)
  - DM weakly couples to SM particles, it can be produced in SM particle annihilations at accelerators
  - Involve **light dark sector mediators** too


### Colliders



### Direct




### Indirect




# Light dark sector

- No evidence of DM at electro-weak scale in experiments
- Light DM with mass of  $M \sim \mathcal{O}(\text{MeV-GeV})$  theoretically well motivated
- **Light dark mediators** involved in the interaction with SM and DM
  - Several signatures depending on the relation between the mediator mass and the DM mass




- Theoretical description through interaction “**portals**”
- They may solve “DM puzzle” and explain observed anomalies [1, 2, 3] like the  $(g - 2)_\mu$

## Portals



Vector	$\epsilon F^{\mu\nu} A'_{\mu\nu}$
Scalar	$k  H ^2  S ^2$
Fermion	$y H L N$
Pseudo-scalar	$\frac{1}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu} a$




[1] Abi et al., *Phys. Rev. Lett.* **126**, 141801 (2021)  
 [2] G. Caria et al. *Phys. Rev. Lett.* **124**, 161803 (2020)  
 [3] R. Aaij et al. *Nature Physics* **18**, 277 (2022)

# The Belle II experiment @ SuperKEKB


See Pavel Pakhlov's talk for details!

- SuperKEKB is a new generation  $B$ -factory  $\rightarrow$  asymmetric  $e^+/e^-$  collider, mainly operated at  $\sqrt{s} = 10.58$  GeV [Y(4S)]
- Belle II is the upgrade of Belle @ KEKB  $\rightarrow$  Hermetic detector with high performances
- $424 \text{ fb}^{-1}$  collected, currently not in data taking
- Well known initial-state condition and clean environment (Low/no pile-up)
- Dedicated low-multiplicity triggers
  - $\rightarrow$  Suppress high-cross-section QED processes without “killing” the signal
  - $\rightarrow$  Precise knowledge of acceptance and efficiencies of the detector required
  - $\rightarrow$  Example: single-photon trigger available in the full collected data set  $\rightarrow$  makes Belle II dataset unique
- Excellent reconstruction capabilities for low multiplicities and missing energy signatures





# Recent dark sector results at Belle II





Belle II detector

# Search for an invisible $Z'$ boson

[1] Shuve et al., *Phys. Rev. D* 89, 113004 (2014)  
 [2] D. Curtin et al., *JHEP* 02 (2015) 157  
 [3] Altmannshofer et al., *JHEP* 106 (2016)

- Massive  $Z'$  boson with a coupling  $g'$  only to leptons with  $\mu$ - and  $\tau$ -lepton numbers ( $L_\mu - L_\tau$  extension of the SM) [1,2,3]
  - It may explain  $(g - 2)_\mu$  anomaly and DM abundance
- Possible decays:
  - $Z' \rightarrow$  invisible ( $\nu\bar{\nu}$  or  $\chi\bar{\chi}$ ),  $Z' \rightarrow \mu\mu$ ,  $Z' \rightarrow \tau\tau$
- $Z' \rightarrow$  invisible ( $Z' \rightarrow \nu\bar{\nu}/\chi\bar{\chi}$ )
  - If light DM  $\chi$  kinematically accessible exists,  $BR(Z' \rightarrow \text{invisible}) = 100\%$
  - Profit from the excellent Belle II capabilities for missing energy signatures
  - Searched for through the process  $e^+ e^- \rightarrow \mu^+ \mu^- Z', Z' \rightarrow \text{inv.}$
  - Signal signature is a narrow peak in the recoil mass of the two final-state muons

$L_\mu - L_\tau$  model  $Z'$  branching ratios in leptons





$$M_{recoil}^2(\mu\mu) = s + M(\mu\mu)^2 - 2\sqrt{s}(E_{\mu^+}^{CMS} + E_{\mu^-}^{CMS})$$



# Search for a $\mu\mu$ -resonance in $e^+e^- \rightarrow \mu^+\mu^-\mu^+\mu^-$

★ New for [EPS-HEP2023](#)

- Four-track final state with at least **three identified as muons**
  - Four-track invariant mass compatible with collision  $\sqrt{s}$
  - No extra energy
- Signal signature is a **narrow peak in the opposite-charge di-muon mass  $M(\mu\mu)$**
- Challenging aggressive suppression of main **SM background  $e^+e^- \rightarrow \mu^+\mu^-\mu^+\mu^-$** 
  - Based on classifiers trained exploiting the features of kinematic distributions in signal events
    - ▶ Presence of a resonance in both **candidate** and **recoil** muon pairs
- Signal extracted through fits to  $M(\mu\mu)$







# Search for a $\mu\mu$ -resonance in $e^+e^- \rightarrow \mu^+\mu^-\mu^+\mu^-$ : results

★ New for [EPS-HEP2023](#)

[1] P. Harris et al., [arxiv-2207.08990 \(2022\)](#)  
 [2] S. Gori et al., [arxiv-2209.04671 \(2022\)](#)


- No significant excess found in  $178 \text{ fb}^{-1}$ 
  - Competitive 90% CL upper limits on the  $g'$  coupling of the  $L_\mu - L_\tau$  model ( $Z'$ ) with *BaBar* ( $> 500 \text{ fb}^{-1}$ ) and Belle ( $> 600 \text{ fb}^{-1}$ ) results
  - First 90% CL upper limits for the muonphilic scalar model from a dedicated search [1, 2]



# Search for a $\tau\tau$ -resonance in $e^+e^- \rightarrow \mu^+\mu^-\tau^+\tau^-$

I. Adachi et al., [arXiv:2306.12294 \(2023\)](https://arxiv.org/abs/2306.12294) - accepted by PRL

- Four-track final state:  $\tau$  decay in  $\tau \rightarrow l\nu\bar{\nu}$ ,  $\tau \rightarrow h\nu\bar{\nu}$
- Signal peaks in the recoil mass of  $\mu^+\mu^-$   $M_{\text{recoil}}(\mu\mu)$
- Challenging background rejection to reduce event contamination with missing energy not associated with signal signature




- Eight classifiers trained on different regions of recoil mass
  - ▶ Based on resonance  $X$  properties (FSR) and  $\tau\tau$  system

- Signal extracted through fit to  $M_{\text{recoil}}(\mu\mu)$  distribution

- Background measured directly on data to minimize impact of not correctly simulated backgrounds

- Smooth background → not problematic





# Search for a $\tau\tau$ -resonance in $e^+e^- \rightarrow \mu^+\mu^-\tau^+\tau^-$ : results

I. Adachi et al., [arXiv:2306.12294 \(2023\)](https://arxiv.org/abs/2306.12294) - accepted by PRL

[1] J. P. Lees et al., [PhysRevLett.125.181801 \(2020\)](https://arxiv.org/abs/2001.06072)

[2] M. Bauer et al., [JHEP09-056 \(2022\)](https://arxiv.org/abs/2205.08609)

- No significant excess found in  $62.8 \text{ fb}^{-1}$ 
  - First limits at 90% CL for a leptophilic dark scalar  $S$  model with  $m_S > 6.5 \text{ GeV}/c^2$  [1]
  - First direct limits at 90% CL for axion-like particle  $ALP \rightarrow \tau\tau$  [2]





# Long-lived spin-0 boson in $b \rightarrow s$ transitions

I. Adachi et al., [arXiv:2306.02830 \(2023\)](https://arxiv.org/abs/2306.02830)

- Search for a **new scalar  $S$**  in  **$B$  meson** decays in  $b \rightarrow s$  transitions
  - ➔  $S$  can mix with SM Higgs boson with mixing angle  $\theta_s$   
→ natural long-lived particle (**LLP**) for small  $\theta_s$
  - ➔ **High performance in LLP vertex reconstruction** are necessary
- **$B$  meson** decays
  - ➔ Eight exclusive “visible” channels reconstructed
  - ➔ Prompt decay of  $K$  or  $K^*$  + **opposite-charged tracks** that make a **displaced vertex**
  - ➔ Backgrounds: combinatorial  $e^+e^- \rightarrow q\bar{q}$ ,  $K_S$  vetoed in  $M_{\pi\pi}$  mass, additional peaking backgrounds suppressed with tighter selections on displaced vertices
- Signal extracted through **fit to the LLP reduced mass, separately for each channel and lifetime**

$$\begin{aligned}
 &B^+ \rightarrow K^+ S \\
 &B^0 \rightarrow [K^{*0} \rightarrow K^+ \pi^-] S \\
 &S \rightarrow ee/\mu\mu/\pi\pi/KK
 \end{aligned}$$






# Long-lived spin-0 boson in $b \rightarrow s$ transitions: results


I. Adachi et al., [arXiv:2306.02830 \(2023\)](https://arxiv.org/abs/2306.02830)

- No significant excess observed in  $189 \text{ fb}^{-1}$ 
  - First model-independent limits at 95% CL on  $BR(B \rightarrow K_S) \cdot BR(S \rightarrow x^+x^-)$
  - First limits on decays to hadrons
- Interpretation as dark scalar  $S$  [1, 2]

[1] [Phys. Rev. D 101, 095006 \(2020\)](https://arxiv.org/abs/2009.09506)  
 [2] [J. Phys. G: Nucl. Part. Phys. 47 010501](https://arxiv.org/abs/1701.01050)






Limits for each channel and lifetime



# $\tau \rightarrow l \alpha$ (invisible) decay

I. Adachi et al., [Phys. Rev. Lett. 130, 181803 \(2023\)](#)

- Charged-Lepton Flavour Violation (LFV) is allowed in various SM extensions  $\rightarrow$  it has never been observed
- $\tau$ -decays in new  $\alpha$  bosons that mediate LFV processes are predicted in different theoretical models [1]
- Search for  $e^+ e^- \rightarrow \tau_{\text{sig}} \tau_{\text{tag}}, \tau_{\text{tag}} \rightarrow 3\pi \nu$
- The presence of neutrinos does not allow to define the reference frame in which  $\tau_{\text{sig}}$  is at rest
- $\rightarrow$  Introduce the approximate  $\tau_{\text{sig}}$  reference frame
- Search for a peak in the normalized energy spectrum of the lepton  $x_l$  (in the approximate  $\tau_{\text{sig}}$  reference frame) over the irriducible SM  $\tau \rightarrow l \bar{\nu} \nu$  background





[1] M. Bauer, et al. [Phys. Rev. Lett. 124, 211803 \(2020\)](#)

# $\tau \rightarrow l \alpha$ (invisible) decay: results

I. Adachi et al., [Phys. Rev. Lett. 130, 181803 \(2023\)](#)


[1] ARGUS Collaboration, [Z. Phys. C 68, 25 \(1995\)](#)

- No excess observed in 62.8 fb<sup>-1</sup>
  - ➔ Limits from 2.2 to 14 times more stringent with respect to the previous existing limits set by ARGUS [1]



# Summary and conclusions

- Belle II has a **unique sensitivity to light dark sector** and progressively will lead its exploration at luminosity frontier
  - ➔ Complementary results to higher energy colliders and beam-dump experiments
  - ➔ World-leading results already published with partial datasets ( $< 424 \text{ fb}^{-1}$ ), updates and new results are coming!
- ▶ Search for an invisible  $Z'$  in  $ee \rightarrow \mu\mu Z'$  [Phys. Rev. Lett. 130, 231801 \(2023\)](#)
- ▶ Search for a resonance decaying to  $\mu\mu$  in  $ee \rightarrow \mu\mu\mu\mu$  events [New for EPS-HEP2023](#)
- ▶ Search for a resonance decaying to  $\tau\tau$  in  $ee \rightarrow \mu\mu\tau\tau$  events [arXiv:2306.12294 \(2023\)](#)
- ▶ Search for a long-lived spin-0 boson in  $b \rightarrow s$  transitions [arXiv:2306.02830 \(2023\)](#)
- ▶ Search for the LFV  $\tau \rightarrow l \alpha$  (invisible) decay [Phys. Rev. Lett. 130, 181803 \(2023\)](#)



Belle II target integrated luminosity is  $50 \text{ ab}^{-1}$

Thank you!







Backup slides

# Belle II perspectives

- Target integrated luminosity: 50  $\text{ab}^{-1}$
- Target peak luminosity:  $6 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$






- 424  $\text{fb}^{-1}$  collected
- **Obtained results are strongly limited by statistics**
  - World-leading results already published with early datasets ( $< 20\%$  of the collected dataset of 424  $\text{fb}^{-1}$ )

- In next years, Belle II will collect 100-times the dataset collected up to now
- ➔ **The best is yet to come!**

# Experiments at B-factories



- Asymmetric  $e^+e^-$  colliders optimized for the production of  $B$  meson pairs, but also  $D$  mesons,  $\tau$  leptons, ...
- Collisions occur at  $\Upsilon(nS)$  resonances
  - ➔ Mainly at  $\Upsilon(4S)$ :  $\sqrt{s} = 10.58$  GeV just above the production threshold of  $B\bar{B}$   
 $BR(\Upsilon(4S) \rightarrow B\bar{B}) > 96\%$
- Asymmetric beam energies: boosted  $B\bar{B}$  pairs, for CP-violation time-dependent measurements
- High peak luminosity  $L > 10^{34} \text{ cm}^{-2}\text{s}^{-1}$




**Belle@KEKB, KEK, Tsukuba (JP)**  
1999–2010,  $\int L dt = 1 \text{ ab}^{-1}$

### First generation of B-factories

**BABAR@PEP-II, SLAC (USA)**  
1999–2008,  $\int L dt = 0.5 \text{ ab}^{-1}$

### Integrated luminosity of B factories





**Tot: 1.5 ab<sup>-1</sup>**


Category	Resonance	Luminosity (fb <sup>-1</sup> )
On resonance	$\Upsilon(5S)$	121
	$\Upsilon(4S)$	711
	$\Upsilon(3S)$	3
	$\Upsilon(2S)$	25
	$\Upsilon(1S)$	6
Off resonance		~100
~550 fb <sup>-1</sup>	On resonance	
	$\Upsilon(4S)$	433
	$\Upsilon(3S)$	30
	$\Upsilon(2S)$	14
	Off resonance	

# SuperKEKB

- New generation of B-factory that provides luminosity to the Belle II experiment
- ➔ Asymmetric beam energies:  $e^-$  (7 GeV) /  $e^+$  (4 GeV)  
Operating mainly at Y(4S), but foreseen runs from Y(2S) to Y(6S)
- ➔ Designed to reach the world highest peak luminosity with the nano-beam scheme




- World record luminosity on December 2021:  $3.8 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- $I(e^-/e^+) = 820/1034 \text{ mA}$  and  $\beta_y^* = 1 \text{ mm}$
- Target peak luminosity:  $6.5 \cdot 10^{35} \text{ cm}^{-2}\text{s}^{-1}$






# Belle II at SuperKEKB

$$\beta\gamma = 0.28$$





- Major upgrade of Belle@KEKB  $\rightarrow$  better resolution, particle identification (PID) and capability to cope with higher background
- Covers more than 90% of the total solid angle

- First collisions during commissioning run on April 26<sup>th</sup> 2018
  - $\rightarrow$   $0.5\ \text{fb}^{-1}$  collected in 2018
- First collisions with the full detector on March 2019
  - $\rightarrow$   $\sim 430\ \text{fb}^{-1}$  collected in 3 years of data taking
- Target integrated luminosity of the Belle II experiment:  **$50\ \text{ab}^{-1}$**  (x30 Belle + BaBar)





# Low-multiplicity triggers

- Two-level trigger
  - Hardware-based Level 1 Trigger (L1): < 30 kHz
  - Software-based High Level Trigger (HLT): < 10 kHz
- Devised specific low-multiplicity trigger lines
  - Suppress high-cross-section QED processes **without “killing” the signal**
  - **Precise knowledge of acceptance and efficiencies of the detector required**
- Examples
  - Single-photon trigger
  - Single-muon trigger
  - Single-track trigger



# Axion-like particles (ALPs)


- GeV-scale ALPs: pseudo-scalar portal mediator between dark sector and Standard Model
- If ALP-photon coupling ( $g_{a\gamma\gamma}$ ) dominates, then  $BR(a \rightarrow \gamma\gamma) \sim 100\%$
- Focus on mass region where ALP decay is prompt and photons can be well **resolved** by Belle II



# Search for an ALP at Belle II


F. Abudinén et al., [Phys. Rev. Lett. 125, 161806 \(2020\)](#)

- Event selection:
  - electromagnetic calorimeter trigger (efficiency  $\sim 100\%$ )
  - three- $\gamma$  invariant mass compatible with collision  $\sqrt{s}$
- Signal signature is a **narrow peak** in  $M_{\gamma\gamma}^2$  or  $M_{\text{recoil}}^2$  (depending on best resolution of signal peak)
- Largest background from  $e^+e^- \rightarrow \gamma\gamma(\gamma)$
- Signal extracted through fit
  - **No excess observed in  $0.445 \text{ fb}^{-1}$**
  - Upper limits at 95% CL on  $g_{a\gamma\gamma}$
  - **World-leading limits for  $m_a \sim 0.5 \text{ GeV}/c^2$**




# Search for a dark Higgs (and dark photon)

- Dark photon  $A'$ 
  - kinetic mixing with SM photon with strength  $\epsilon$  [1]
  - mass produced by the Higgs mechanism involving a dark Higgs boson [2]
- Dark higgs  $h'$ 
  - couples to  $A'$  with  $\alpha_D$
  - does not mix with SM Higgs
- Both  $A'$  and  $h'$  can be produced at  $e^+e^-$  colliders through the dark higgsstrahlung process
  - $e^+e^- \rightarrow A'^* \rightarrow A' h'$
- Different signatures depending on  $h'$  mass
  - $M_{h'} > M_{A'}$ : prompt decay  $h' \rightarrow A'A'$ , up to 6 tracks in the final state. Investigated by [BaBar\(2012\)](#) and [Belle\(2015\)](#)
  - $M_{h'} < M_{A'}$ :  $h'$  is long-lived, thus invisible. Investigated by [KLOE\(2015\)](#)
- Belle II focuses on the invisible  $h'$



[1] P. Fayet, [Nucl. Phys. B 187, 184 \(1981\)](#)

[2] Batell et al., [Phys. Rev. D 79, 115008 \(2009\)](#)








# Dark higgsstrahlung at Belle II

F. Abudinén et al., *Phys. Rev. Lett.* **130**, 071804 (2023)


- $e^+e^- \rightarrow A'h', A' \rightarrow \mu\mu, h' \rightarrow \text{invisible}$
- Same final state as for the invisible  $Z'$ , similar backgrounds:  
 $e^+e^- \rightarrow \tau^+\tau^- (\gamma), e^+e^- \rightarrow \mu^+\mu^- (\gamma), e^+e^- \rightarrow e^+e^-\mu^+\mu^-$
- Signal signature is a 2D peak in the recoil mass vs the dimuon mass
- Event selection
  - ➔ Two reconstructed muons,  $p_{T^\mu} > 0.1 \text{ GeV}/c$
  - ➔ Recoil momentum in the ECL barrel, no nearby photon
  - ➔ Cut on dimuon helicity angle  
 ➔ efficiently suppress background
- Signal extraction through 2D fit in  $M_{\text{recoil}}$  vs  $M_{\mu\mu}$  plane in elliptical windows





# Dark higgsstrahlung at Belle II: results

F. Abudinén et al., *Phys. Rev. Lett.* **130**, 071804 (2023)

- **No significant excess in 8.34 fb<sup>-1</sup>**
  - 90% CL upper limits and world leading limits for  $1.65 < M_{A'} < 10.51 \text{ GeV}/c^2$



Cross section [fb]



# Search for dark photon $A'$

[1] P. Fayet, [Phys. Lett. B 95, 285 \(1980\)](#)  
 [2] P. Fayet, [Nucl. Phys. B 187, 184 \(1981\)](#)

- U(1) extension of the SM
- New massive vector gauge boson,  $A'$ , with a coupling to the Standard Model photon through the kinetic mixing mechanism, with strength  $\epsilon$  [1,2]

$$\mathcal{L}_{int} = e \underbrace{\epsilon}_{\text{Interaction strength}} \underbrace{A'_\mu}_{\text{Dark photon field}} \underbrace{J_{em}^\mu}_{\text{Electromagnetic current}}$$


- This gauge boson can be produced at  $e^+e^-$  colliders through different processes:

- direct production:  $e^+e^- \rightarrow \gamma_{ISR} A'$
- meson decays:  $\pi^0 \rightarrow A' \gamma$
- dark higgsstrahlung:  $e^+e^- \rightarrow A'^* \rightarrow A' h'$

- **Direct production with ISR particularly interesting:**  $e^+e^- \rightarrow \gamma_{ISR} A'$


- Two basic scenarios depending on dark photon mass:

- $M_{A'} > 2m_\chi$ : invisible decay  $A' \rightarrow \chi\bar{\chi}$
- $M_{A'} < 2m_\chi$ : visible decay in Standard Model particles




# Invisible dark photon

- $e^+e^- \rightarrow \gamma_{ISR} A' (A' \rightarrow inv.)$ 
  - ➔ Single photon search: single photon trigger needed, present in the full Belle II dataset




[1] Belle II Physics Book, PTEP 2019 12 (2019)

[2] Less et al, Phys. Rev. Lett. 119, 131804 (2017)





Background simulation assuming 20 fb<sup>-1</sup>



- Belle II expected to perform better than BABAR [2]:
  - smaller boost: larger acceptance
  - muon detector veto: reject events with a photon undetected in the calorimeter (efficiency currently under study)
  - better calorimeter hermeticity

# Visible dark photon

- BABAR [1]
  - Full dataset of  $514 \text{ fb}^{-1}$
  - Dark photon visible decay in  $e^+e^-$  and  $\mu^+\mu^-$  final states
  - Signal signature, bump in the dilepton invariant mass
  - Background: QED processes  $e^+e^- \rightarrow e^+e^-(\gamma)$ ,  $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$  and resonant backgrounds from  $J/\psi$ ,  $\psi(2S)$  etc. (vetoed)
  - Upper limits at 90% CL kinetic mixing strength  $\varepsilon$  at level of  $O(10^{-3})$ :
- LHCb [2]
  - Best limits in the mass range 200 -700 MeV
- Belle II is expected to achieve the leading sensitivity [3]
  - Search in preparation



[1] J.P. Lees et al, *Phys. Rev. Lett.* 113, 201801 (2014)

[2] R. Aaij et al, *PhysRevLett.* 124.041801 (2020)

[3] E. Kou et al, *Prog Theor Exp Phys* (2019)



# Search for a $\tau\tau$ -resonance in $e^+e^- \rightarrow \mu^+\mu^-\tau^+\tau^-: Z'$

- No significant excess found in  $62.8 \text{ fb}^{-1}$

→ 90% CL upper limits on the  $g'$  coupling of the  $L_\mu - L_\tau$  model ( $Z'$ )

