

*New Results on Beauty, Charm,
and Tau from Belle II*

*Soeren Prell (Iowa State University)
14th Conference on the Intersections of
Particle and Nuclear Physics (CIPANP 2022)*

Orlando, Florida



August 29 – September 4, 2022

On behalf of the Belle II Collaboration



Belle II & SuperKEKB Accelerator


- *Belle II is a multipurpose experiment at the SuperKEKB collider located at KEK (Tsukuba, Japan)*
 - *Asymmetric-energy e^+ (4 GeV) e^- (7 GeV) collider with E_{CM} near the $\Upsilon(4S)$ resonance (~ 10.6 GeV)*
- *Aims to collect a 50 ab^{-1} data sample ($50 \times$ Belle)*
- *Final design instantaneous luminosity of $6.5 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ ($30 \times$ that of KEKB) by*
 - *reducing beam size by factor 20 (“nano beams”)*
 - *increasing beam current by factor 1.5*



factor 20


factor 1.5

Factor ~ 30 in the luminosity





Belle II Detector





Belle II, a Super Heavy Flavor Factory

Belle II is ...


- a Super **B** Factory:
 $1.1 \times 10^9 B\bar{B}$ pairs per ab^{-1}
- a Super **Charm** Factory:
 $1.3 \times 10^9 c\bar{c}$ pairs per ab^{-1}
- a Super **τ** Factory:
 $0.9 \times 10^9 \tau^+\tau^-$ pairs per ab^{-1}

... and in addition, the clean e^+e^- environment allows the study/search of

- Charmonium & bottomonium (SM & exotic X,Y,Z)
- Tetra- and penta-quarks
- Dark particles (dark γ /Higgs, ALPs, LLPs), ...



Belle II Luminosity



*Belle II has recorded
428 fb⁻¹ in 2019-2022*

- *On-Υ(4S): 363 fb⁻¹*
- *Below-Υ(4S): 42 fb⁻¹*
- *Unique 10.75 GeV scan:
19 fb⁻¹*

*Today's results are based
on up to 190 fb⁻¹ On-Υ(4S)*

*Integrated luminosities
On-Υ(4S) of B factories
for comparison:
Belle (711 fb⁻¹)
BaBar (424 fb⁻¹)*

*Many Belle II results are starting to become statistically competitive,
some measurements are already world's best !*







B Meson Reconstruction Techniques

Exploit kinematics of $e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$ for signal selection

$$M_{bc} = \sqrt{(E_{beam}^*)^2 - (p_B^*)^2}$$

$$\Delta E = E_B^* - E_{beam}^*$$


Event shape variables




Full Event Interpretation

Comput. Softw. Big Sci. 3, 6 (2019)



- *Reconstruct one B (B_{tag}) fully w/ multivar. classifiers (hadr. and SL final states)*
 - *>200 BDTs reconstr. $O(10k)$ decay chains*
 - *Samples dominated by large-BF, low-background modes*
- *Properties of B_{sig} (e.g. momentum) and invisible daughters (e.g. “missing mass”) can be calculated with B_{tag} momentum*
- *$\sim(1.5-2) \times$ better perform. than at Belle*
 - *Typical for values hadronic B_{tag} : $\epsilon(B_{tag}^+) \sim 0.5\%$, purity $\sim 10\%$*




Example:



Hadronic decay of B_{tag}



Signal decay with invisible neutrino



Measurements of quark mixing parameters

Amplitude for charged current quark transition $q_i \rightarrow W q_j$ is proportional to CKM matrix element V_{ij}


- BFs ($\propto |V_{ij}|^2$) \rightarrow magnitudes
- CP asymmetries (arising from interference of 2 amplitudes) \rightarrow (complex) phases

$$V_{\text{CKM}} \equiv V_L^u V_L^{d\dagger} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$

Belle II can measure magnitudes of 7 of the 9 matrix elements and the weak phase

In the SM, V_{CKM} is a unitary 3×3 matrix: measurements of Unitarity Triangle sides and angles must be consistent !!!

$$V_{ub}^* V_{ud} + V_{cb}^* V_{cd} + V_{tb}^* V_{td} = 0$$





$$\begin{aligned} \phi_1 &= \beta \equiv \arg [-V_{cd} V_{cb}^* / V_{td} V_{tb}^*] \\ \phi_2 &= \alpha \equiv \arg [-V_{td} V_{tb}^* / V_{ud} V_{ub}^*] \\ \phi_3 &= \gamma \equiv \arg [-V_{ud} V_{ub}^* / V_{cd} V_{cb}^*] \end{aligned}$$

V_{ub} and V_{cb}

$|V_{ub}|$ and $|V_{cb}|$ are precisely measured with semileptonic B decays

Exclusive: $B \rightarrow \pi/\rho \ell \nu$, $B \rightarrow D(*) \ell \nu$ etc.

Inclusive: $B \rightarrow X_u \ell \nu$, $B \rightarrow X_c \ell \nu$




Parameter	Exclusive	Inclusive
$ V_{cb} \times 10^{-3}$	39.10 ± 0.50	42.19 ± 0.78
$ V_{ub} \times 10^{-3}$	3.51 ± 0.12	4.19 ± 0.12

HFLAV, [arXiv:2206.07501](https://arxiv.org/abs/2206.07501)


discrepancy between inclusive and exclusive

V_{cb} from untagged $B \rightarrow D \ell \nu$ sample



θ_{BY} is angle between B & Dl system

V_{ub} from tagged $B \rightarrow \pi \ell \nu$ sample



4 new exclusive V_{xb} results from Belle II:

$ V_{xb} $	Signal B (B_{sig}) decay	Other B (B_{tag}) decay	Latest result
$ V_{cb} $	$B_{sig} \rightarrow D\ell\nu$ ($\ell=e,\mu$)	untagged	ICHEP 2022
$ V_{cb} $	$B^0_{sig} \rightarrow D^*\ell\nu$ ($\ell=e,\mu$)	hadronically tagged	Moriond 2022
$ V_{ub} $	$B^0_{sig} \rightarrow \pi\ell\nu$ ($\ell=e,\mu$)	untagged	ICHEP 2022
$ V_{ub} $	$B_{sig} \rightarrow \pi e\nu$	hadronically tagged	Moriond 2022

$|V_{xb}| \times 10^3$



$\eta_{EW}|V_{cb}| = 38.53 \pm 1.15$

$\eta_{EW}|V_{cb}| = 38.2 \pm 2.8$

$|V_{ub}| = 3.54 \pm 0.25$

$|V_{ub}| = 3.88 \pm 0.45$

η_{EW} = EW correction factor



Belle II V_{xb} results are consistent with previous measurements, with precision approaching those of prev. results

Excl. vs incl. discrepancy may be due to unaccounted non-perturbative effects.

Measurements of SL decay kinematics (Belle II, arxiv.org:2205.06372) may help resolve the issue.

Meier, HF
Sat 15:30 &

Time-dependent asymmetries in B decays

- *TD measurements of B decays at $\Upsilon(4S)$ were pioneered by BABAR and Belle*
 - *Need good B flavor tagging and $\Delta t = t_{\text{sig}} - t_{\text{tag}}$ measurement*
- *$B\bar{B}$ mixing freq. Δm_d determined from TD of $\Upsilon(4S) \rightarrow B\bar{B}$ or $BB, \bar{B}\bar{B}$*



$$\mathcal{A}(\Delta t) = \frac{N_{B\bar{B}} - N_{BB, \bar{B}\bar{B}}}{N_{B\bar{B}} + N_{BB, \bar{B}\bar{B}}} = \cos(\Delta m_d \Delta t) (1 - 2w) \otimes R(\Delta t)$$

- *New Belle II measurements of B lifetime and mixing frequency*

$$\tau_{B^0} = 1.499 \pm 0.013 \pm 0.008 \text{ ps}$$

$$\Delta m_d = 0.516 \pm 0.008 \pm 0.005 \text{ ps}^{-1}$$


- *Measurements consistent with WAs*
- *O(1%) precision in $\tau(B^0)$ and Δm_d demonstrates Belle II's excellent flavor tagging and vertexing performance*



Measurement of CP asymmetry $\sin 2\phi_1$

$$A^{\text{raw}}(\Delta t) = \frac{N(\bar{B}^0 \rightarrow f_{CP}) - N(B^0 \rightarrow f_{CP})}{N(\bar{B}^0 \rightarrow f_{CP}) + N(B^0 \rightarrow f_{CP})}(\Delta t) = \underbrace{A_{CP}}_{\text{direct CP asymmetry}} \cos(\Delta m_d \Delta t) + \underbrace{S_{CP}}_{\text{mixing-induced CP asymmetry}} \sin(\Delta m_d \Delta t)$$

- Expect $S_{CP} = \sin 2\phi_1$ for tree amplitude $b \rightarrow c\bar{c}s$ decays
- New physics could provide CP contribution in penguin decays




- New Belle II $\sin 2\phi_1$ measurement with golden mode $B \rightarrow J/\psi K_S^0$

$$S_{CP} = 0.720 \pm 0.062 \text{ (stat)} \pm 0.016 \text{ (syst)}$$

$$A_{CP} = 0.094 \pm 0.044 \text{ (stat)} \begin{matrix} +0.042 \\ -0.017 \end{matrix} \text{ (syst)}$$

- 9% measurement is statistically dominated
- Consistent with WA



Time-dependent CPV in B penguins



ICHEP 2022

- Measure S_{CP} in penguin decay $B \rightarrow 3K_S^0$
- Result are consistent with SM predictions:
 - $A_{CP} \sim 0$ and $S_{CP} \sim -\sin 2\phi_1$

$$S_{CP} = -1.86_{-0.46}^{+0.91} \text{ (stat)} \pm 0.09 \text{ (syst)}$$

$$A_{CP} = -0.22_{-0.27}^{+0.30} \text{ (stat)} \pm 0.04 \text{ (syst)}$$

- Technically complicated measurement with no tracks from B_{sig} decay vertex
- Small inner radius of PXD ensures most $K_S^0 \rightarrow \pi^+\pi^-$ daughter tracks have pixel hit info



$K\pi$ Puzzle

- Unexpected large difference between CP asymmetries $A_{K^+\pi^-}^{CP}$ and $A_{K^+\pi^0}^{CP}$ in $B \rightarrow K\pi$ decays dominated by hadronic penguin amplitudes
- Isospin sum rule tests if discrepancy from sub-leading SM amplitudes

$$I_{K\pi} = \mathcal{A}_{K^+\pi^-}^{CP} + \mathcal{A}_{K^0\pi^+}^{CP} \frac{\mathcal{B}_{K^0\pi^+}}{\mathcal{B}_{K^+\pi^-}} \frac{\tau_{B^0}}{\tau_{B^+}} - 2\mathcal{A}_{K^+\pi^0}^{CP} \frac{\mathcal{B}_{K^+\pi^0}}{\mathcal{B}_{K^+\pi^-}} \frac{\tau_{B^0}}{\tau_{B^+}} - 2\mathcal{A}_{K^0\pi^0}^{CP} \frac{\mathcal{B}_{K^0\pi^0}}{\mathcal{B}_{K^+\pi^-}} \approx 0$$


- Current precision (13%) limited by $A_{K^0\pi^0}^{CP}$.
- Only Belle II can measure all of these !

New Belle II measurements:

$$\begin{aligned} A_{CP}^{K^+\pi^0} &= 0.014 \pm 0.047 \pm 0.010 \\ B_{K^+\pi^0} &= (14.30 \pm 0.69 \pm 0.79) \times 10^{-6} \\ A_{CP}^{K^0\pi^0} &= -0.41_{-0.32}^{+0.30} \pm 0.09 \\ B_{K^0\pi^0} &= (11.0 \pm 1.2 \pm 1.0) \times 10^{-6} \end{aligned}$$

ICHEP 2022


Previous Belle II results with 63 fb^{-1} : $K^+\pi^-$ and $K^0\pi^+$ (arXiv:2106.03766), $K^0\pi^0$ (arXiv:2104.14871), and $K^+\pi^0$ (arXiv:2105.04111),



Measurement of ϕ_2 from $B \rightarrow \pi\pi$ and $B \rightarrow \rho\rho$

CKM angle ϕ_2 determined through BFs and CP asymmetries for sets of $b \rightarrow u$ dominated $B \rightarrow \pi\pi$ and $B \rightarrow \rho\rho$ decays

- Isospin decomposition is necessary to account for penguin pollution



New Belle II $B \rightarrow \pi\pi$ measurements:



$$A_{CP}^{\pi^+ \pi^0} = -0.085 \pm 0.085 \pm 0.019$$

$$B_{\pi^+ \pi^0} = (6.12 \pm 0.53 \pm 0.53) \times 10^{-6}$$

$$A_{CP}^{\pi^0 \pi^0} = 0.14 \pm 0.46 \pm 0.07$$

$$B_{\pi^0 \pi^0} = (1.27 \pm 0.25 \pm 0.17) \times 10^{-6}$$

WA: $A_{CP}^{\pi^0 \pi^0} = 0.33 \pm 0.22$, $B_{\pi^0 \pi^0} = (1.59 \pm 0.26) \times 10^{-6}$



Although $\rho\rho$ is a VV final state, similar isospin analysis as in $\pi\pi$ possible since only longitudinal amplitude dominant

New Belle II $B \rightarrow \rho\rho$ measurements:

$$B_{\rho^+\rho^-} = (26.7 \pm 2.8 \pm 2.8) \times 10^{-6}$$

$$f_L^{\rho^+\rho^-} = 0.956 \pm 0.035 \pm 0.033$$

WA: $B_{\rho^+\rho^-} = (27.7 \pm 1.9) \times 10^{-6}$



$$A_{CP}^{\rho^+\rho^0} = -0.069 \pm 0.068 \pm 0.060$$

$$B_{\rho^+\rho^0} = (23.2_{-2.1}^{+2.2} \pm 2.7) \times 10^{-6}$$

$$f_L^{\rho^+\rho^0} = 0.943_{-0.033}^{+0.035} \pm 0.027$$


WA: $A_{CP}^{\rho^+\rho^0} = -0.05 \pm 0.05, B_{\rho^+\rho^0} = (24.0 \pm 1.9) \times 10^{-6}$

Previous Belle II results with 63 fb^{-1} : $\pi^+\pi^-$ (arXiv:2106.03766), $\pi^0\pi^0$ (arXiv:2107.02373), and $\rho^+\rho^0$ (arXiv:2206.12362),



Measurement of ϕ_3 with $B^\pm \rightarrow D(K_S^0 h^+ h^-)K^\pm$

Measure ϕ_3 through interference of $b \rightarrow c$ and $b \rightarrow u$ amplitudes in bins of $D/\bar{D} \rightarrow K_S^0 h^+ h^-$ Dalitz plots





Strong phase in D decay from CLEO and BESIII used as external input

$$\phi_3 = (78.4 \pm 11.4 \pm 0.05 \pm 1.0)^\circ$$

WA: $\phi_3 = (65.9^{+3.3}_{-3.5})^\circ$

First joint Belle (711 fb^{-1}) and Belle II (128 fb^{-1}) analysis !



Belle: $N(K_S^0 \pi \pi) = 1467 \pm 53$, $N(K_S^0 KK) = 194 \pm 17$



Belle II: $N(K_S^0 \pi \pi) = 280 \pm 21$, $N(K_S^0 KK) = 34 \pm 7$

Radiative and EW Penguin B Decays

- Flavor-changing neutral currents: in SM due to $b \rightarrow s$ transitions at one-loop level
 - Sensitive to New Physics particles in the loop
- BF ratios, asymmetries and angular observables can be precisely predicted in SM
- New Belle II incl. $BR(b \rightarrow s \gamma)$ measurement
 - Apply cut-off due to large background at low $E(\gamma)$







E_γ^B threshold, GeV	$\mathcal{B}(B \rightarrow X_s \gamma)(10^{-4})$
1.8	3.54 ± 0.78 (stat.) ± 0.83 (syst.)
2.0	3.06 ± 0.56 (stat.) ± 0.47 (syst.)

SM prediction for $E_\gamma^B > 1.6$ GeV: $(3.40 \pm 0.17) \times 10^{-4}$ [JHEP06(2020)175]

$B \rightarrow K^* l^+ l^-$ and $B \rightarrow K \nu \bar{\nu}$

Martel, HI
Thu 16:30

New measurements of $BR(B \rightarrow K^* l^+ l^-)$ and $BR(B \rightarrow K \nu \bar{\nu})$ (fully-incl.)



$$\text{BF}(B^+ \rightarrow K^+ \nu \bar{\nu}) < 4.1 \times 10^{-5}$$

Fully-inclusive method improves sensitivity significantly over previous measurements! 19

Moriond 2022

Decay	Belle II (10^{-6})	PDG (10^{-6})
$B \rightarrow K^* e^+ e^-$	$1.42 \pm 0.48 \pm 0.09$	1.19 ± 0.20
$B \rightarrow K^* \mu^+ \mu^-$	$1.19 \pm 0.31^{+0.08}_{-0.07}$	1.06 ± 0.09

LFU in $B \rightarrow D^{(*)} l \nu$

Hara, HI
Sat 14:00

- Tensions observed recently in excl. semi-leptonic BF ratios $R(D^{(*)}) = BR(B \rightarrow D^{(*)} \tau \nu) / BR(B \rightarrow D^{(*)} l \nu)$

- Predictions for incl. $R(X)$:

$$R(X_{c,\tau/l})_{SM} = 0.223 \pm 0.004$$

PRD 92 (2015) 054018



$$R(X_{c,e/\mu})_{SM} = 1.006 \pm 0.001$$

Vos & Rahimi, in progress

- Since incl. measurements are hard, esp. with τ , measure $R(X_{e/\mu})$ first

$$R(X_{c,e/\mu}) = 1.003 \pm 0.010 \pm 0.020$$

Most precise LFU test with semi-leptonic B decays to date !



Charm meson lifetimes


- First D^0 and D^+ lifetime measurements in 2 decades

Belle II, PRL 127 (2021) 021801



$$\tau(D^0) = 410.5 \pm 1.1 \text{ (stat)} \pm 0.8 \text{ (syst)} \text{ fs}$$

$$\tau(D^+) = 1030.4 \pm 4.7 \text{ (stat)} \pm 3.1 \text{ (syst)} \text{ fs}$$

- Belle II results are more precise than and consistent with previous measurements



0.5% precision (incl. syst.) demonstrates excellent performance and understanding of Belle II vertex detector



Charm baryon lifetimes

- Recent LHCb Λ_c^+ and Ξ_c^0 lifetime measurements changed order of charm baryon lifetimes


- New Belle II results:

$$\begin{aligned} \tau(\Lambda_c^+) &= 203.2 \pm 0.9 \text{ (stat)} \pm 0.8 \text{ (syst)} \text{ fs} \\ \tau(\Omega_c^0) &= 243 \pm 48 \text{ (stat)} \pm 11 \text{ (syst)} \text{ fs} \end{aligned}$$





- Most precise Λ_c^+ lifetime measurement

- Confirms that Ω_c^0 is not shortest-lived singly-charmed baryon

- Consistent with LHCb results
- Inconsistent with pre-LHCb world average by 3.4sigma




arXiv: 2206.15227



LFV searches in τ decays




Many new physics models predict cLFV at 10^{-7} - 10^{-10}



[2207.06307](https://arxiv.org/abs/2207.06307)

$\tau \rightarrow e/\mu + \alpha$ (*invisible*)

- *Invisible particle occur in many NP models (e.g. light ALPs)*
- *Previous best upper limits for $0.1 < M_\alpha < 1.6 \text{ GeV}$ from ARGUS (Z. Phys. C68 (1995) 25)*
- *Compare $\tau \rightarrow e/\mu + \alpha$ rate with SM $\tau \rightarrow e/\mu \nu \bar{\nu}$ prediction*
 - *Improved 95% C.L. limits set for $BR(\tau \rightarrow e/\mu + \alpha)/BR(\tau \rightarrow e/\mu \nu \bar{\nu})$ for $M_\alpha < 1.6 \text{ GeV}$*



Plots and limits for $\tau \rightarrow e\alpha$ are similar !

Conclusions

- *SuperKEKB is delivering $e^+ e^-$ collision data at world-record luminosity*
 - *Expect to improve a factor of 6 during next running period (will start in Fall 2023)*
- *Belle II detector demonstrates excellent performance*
 - *Esp. in incl. reco., neutrals (γ , π^0) & vertex measurements*
- *Belle II is a Super Flavor Factory, already producing many results with first 190 fb^{-1} (of 424 fb^{-1} recorded)*
 - *New B, Charm, and τ physics results are at precisions comparable to those of BABAR and Belle*
 - *Similarly, many new and unique results on dark sector searches & heavy quarkonium (not covered in this talk)*