

Recent results from Belle II

Qi-Dong Zhou
(IAR/KMI, Nagoya Univ.)

On behalf of Belle II Collaboration

May 23 - 29, 2022

21st International Symposium on Very High Energy Cosmic Ray Interactions
ISVHECRI 2022 (Virtual)




New physics search at Belle II




New particle



Energy frontier : direct search



Luminosity frontier : indirect search





- Indirect search for New Physics (NP) in quantum effect
 - Sensitivity of NP detection up to **200 TeV** for loop diagram (depending on the NP coupling constant)
- Standard Model suppressed or forbidden decays
- Test lepton flavor universality and the lepton flavor violations
- Dark sector search, etc.

arXiv:1309.2293

Luminosity frontier: SuperKEKB/Belle II


Asymmetric e^+e^- collider operating at a center of mass energy of the $\gamma(4S)$ resonance



Nano beam scheme

Belle

Belle II



- Squeeze the beam $\sigma_y^* \sim 50 \text{ nm}$
- Large crossing angle

World's highest instantaneous luminosity:

$$\mathcal{L} = 4.14 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$$

KEKB record: $2.1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

The Belle II detector

Vertex detector (VXD)

Inner 2 layers: pixel detector (PXD)
Outer 4 layers: strip sensor (SVD)

e^- (7GeV)

Central Drift Chamber (CDC)


He (50%), C₂H₆ (50%), small
cells, long lever arm

ElectroMagnetic Calorimeter (ECL)

Barrel: CsI(Tl) + waveform sampling
Endcap: pure CsI + waveform
sampling

Level-1 trigger :CDC+ECL+TOP+KLM

DAQ: Maximum 30 kHz L1 trigger



Particle Identification

Barrel: Time-Of-Propagation counters
(TOP)
Forward: Aerogel RICH (ARICH)

e^+ (4GeV)

K_L/μ detector (KLM)


Outer barrel: Resistive Plate Counter
(RPC)
Endcap/inner barrel: Scintillator

Operation status and integrated luminosity

- Belle II operation under COVID-19

Belle II data taking efficiency ~90%

- ~380 fb⁻¹ till now
- Belle: 1 ab⁻¹
- Long shutdown (LS) 1 starts from summer 2022 to autumn 2023 to replace VXD
- LS2 is under discussion for machine improvements on the time frame of 2026-27



CKM matrix and unitarity triangle (UT)

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A^2\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4)$$

Complex phase cause CP violation


$$V^\dagger V = 1 \rightarrow \text{b arrow } \mathbf{d} \text{ column} \rightarrow$$

$$\frac{V_{ub}^* V_{ud}}{\lambda^3 \cdot 1} + \frac{V_{cb}^* V_{cd}}{\lambda^2 \cdot \lambda} + \frac{V_{tb}^* V_{td}}{1 \cdot \lambda^3} = 0$$

$$\phi_1 = \arg \left(-\frac{V_{cd} V_{cb}^*}{V_{td} V_{tb}^*} \right)$$


$$\phi_2 = \arg \left(-\frac{V_{td} V_{tb}^*}{V_{ud} V_{ub}^*} \right)$$

$$\phi_3 = \arg \left(-\frac{V_{ud} V_{ub}^*}{V_{cd} V_{cb}^*} \right)$$






- A triangle on the complex plane
- Normalization by $\bar{\rho} = \rho(1 - \frac{\lambda^2}{2})$
- $\bar{\eta} = \eta(1 - \frac{\lambda^2}{2})$
- Comprehensive test (only Belle II)
 - Measure **all sides and angles**
 - Search NP in mixing (**tree, loop**) by precise measurement of UT

$|V_{cb}|, |V_{ub}|$ measurement through semileptonic B decays



Status of $|V_{ub}|$ and $|V_{cb}|$ determinations

Semi-leptonic decay:





$|V_{ub}|$ and $|V_{cb}|$ longstanding discrepancy btw inclusive and exclusive measurements

Tag side reconstruction at Belle II

- Hadronic tag : Full Event Interpretation trained 200 BDTs to reconstruct ~10000 B decay chains
 - $\varepsilon=0.47\%$ for B^\pm
 - $\varepsilon=0.29\%$ for B^0


[arXiv:2008.06096](https://arxiv.org/abs/2008.06096)




$$m_{bc} = \sqrt{(E_{\text{beam}}^*)^2 - (p_B^*)^2}$$

Measuring $|V_{ub}|$ from $B^0 \rightarrow \pi^- e \nu$

$$\frac{d\mathcal{B}}{dq^2}(B \rightarrow \pi \ell \nu) \propto |V_{ub}|^2 f_+^2(q^2)$$



$$M_{\text{miss}}^2 = (p_{e^+ e^-} - p_{B_{\text{tag}}} - p_e - p_\pi)^2$$





$$q^2 = m_{\ell\nu}^2 = (p_{e^+ e^-} - p_{B_{\text{tag}}} - p_\pi)^2$$

Combined fit of $B^0 \rightarrow \pi^- e^+ \nu$ and $B^+ \rightarrow \pi^0 e^+ \nu$

$$|V_{ub}| = (3.88 \pm 0.45) \times 10^{-3}$$

$$|V_{ub}| = (3.67 \pm 0.15) \times 10^{-3} \text{ (PDG)}$$

Measurement of $B \rightarrow D^* l \bar{\nu}$ for $|V_{cb}|$



$$M_{\text{miss}}^2 = (\cancel{p}_{e^+ e^-} - \cancel{p}_{B_{\text{tag}}} - \cancel{p}_\ell - \cancel{p}_{D^*})^2$$

$$\frac{d\Gamma}{d\omega} \propto |V_{cb}|^2 |\mathcal{F}(\omega)|^2$$

$\mathcal{F}^2(w)$: Form factor determination rely heavily on $w = 1$ (zero recoil), using CLN parameterization, NP B530, 153 (1998)

$$w = \frac{m_B^2 - m_{D^*}^2 - q^2}{2m_B m_{D^*}}$$

w = 1


w = 1.5

$|V_{cb}| = (37.9 \pm 2.7) \times 10^{-3}$


$|V_{cb}| = (39.5 \pm 0.9) \times 10^{-3}$ (PDG)

Measurement of $B \rightarrow X_c l \bar{\nu}$ for $|V_{cb}|$

Hadronic mass moments of
inclusive $B \rightarrow X_c l \bar{\nu}$ with hadronic tag





Publication in preparation



q^2 moments as a function of q^2 momentum threshold


- A new method proposed in [JHEP02 \(2019\)177](#) to extract $|V_{cb}|$ from q^2 moments
- $B \rightarrow X_c l \bar{\nu}$ decay width is expressed with HQE (heavy-quark expansion) parameters
- This method reduce HQE parameters from 13 to 8
- Global fit for inclusive $|V_{cb}|$ in the future

First Belle II $|V_{ub}|$ and $|V_{cb}|$ results



- These are the first Belle II tagged measurements of $|V_{ub}|$ and $|V_{cb}|$ are still statistically limited
 - More precise measurements with larger dataset
 - Higher precision with untagged measurement as the efficiency is 20-30%

$\phi_1/\beta, \phi_2/\alpha, \phi_3/\gamma$ measurements




Time dependent CPV - Flavor tagging

Time dependent CP-Violation measurement:

- Precise measurement of Δt
- B flavor tagger

$$A_{CP} = \frac{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) - \Gamma(B^0(t) \rightarrow f_{CP})}{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) + \Gamma(B^0(t) \rightarrow f_{CP})} = S_f \sin(\Delta mt) + A_f \cos(\Delta mt)$$




S_f : indirect (Time dependent) CPV parameter

A_f : direct CP violation parameter

Δm : the oscillation frequency

$S_f = -\xi \sin(2\phi_I)$ with $B^0 \rightarrow J/\psi K_s$




Effective flavor tagging efficiency:

- Belle II : (30.0 ± 1.3)%
- Belle : (30.1 ± 0.4)%

EPJ C(2022) 82, 283


B^0 Lifetime and mixing frequency



$$\tau_{B^0} = 1.499 \pm 0.013 \text{ (stat.)} \pm 0.008 \text{ (syst.) ps}$$

$$\tau_{B^0} = 1.519 \pm 0.004 \text{ ps (PDG)}$$

$$\Delta m = 0.516 \pm 0.008 \text{ (stat.)} \pm 0.005 \text{ (syst.) ps}^{-1}$$






- Result compatible with world average
- Similar uncertainty as Belle, BaBar results
 - $B \rightarrow D^*/\nu$ to be included
- Belle II ready for time dependent analysis
- Next step $\sin(2\phi_1)$ measurement

ϕ_2/α measurement ($B^+ \rightarrow \rho^+ \rho^0$)

Constraint for ϕ_2 using combination of $B \rightarrow \rho\rho$ ($\rho^+\rho^-$, $\rho^\pm\rho^0$, $\rho^0\rho^0$) decays

- Longitudinal polarization fraction f_L
- Asymmetry in rate $B^+ \rightarrow \rho^+ \rho^0$ vs $B^- \rightarrow \rho^- \rho^0$
 - Direct CP-violation from interference between tree and penguin diagram




$$A_{\text{CP}} = -0.069 \pm 0.068 \text{ (stat.)} \pm 0.060 \text{ (syst.)}$$

$$\mathcal{B}(B^+ \rightarrow \rho^+ \rho^0) = (23.2^{+2.2}_{-2.1} \text{ (stat.)} \pm 2.7 \text{ (syst.)}) \times 10^{-6}$$


$$f_L = 0.943^{+0.035}_{-0.033} \text{ (stat.)} \pm 0.027 \text{ (syst.)}$$

World average: $A_{\text{CP}} = -0.05 \pm 0.05$


ϕ_3/γ measurement



Favored



Color suppressed





- Interference between $b \rightarrow c$ and $b \rightarrow u$ (tree level)

$$\frac{A^{suppr.}(B^- \rightarrow \bar{D}_0 K^-)}{A^{favor.}(B^- \rightarrow D_0 K^-)} = r_B e^{i(\delta_B - \phi_3)}$$

r_B : ratio of amplitude

δ_B : strong phase difference



First combined Belle (711 fb^{-1}) and Belle II (128 fb^{-1}) analysis

$\delta_B [\circ]$	$124.8 \pm 12.9 \text{ (stat.)} \pm 0.5 \text{ (syst.)} \pm 1.7 \text{ (ext.)}$
r_B	$0.123 \pm 0.024 \text{ (stat.)} \pm 0.001 \text{ (syst.)} \pm 0.002 \text{ (ext.)}$
$\gamma [\circ]$	$78.4 \pm 11.4 \text{ (stat.)} \pm 0.5 \text{ (syst.)} \pm 1.0 \text{ (ext.)}$


- Expect $< 3^\circ$ uncertainty with 10 ab^{-1}
- Will still statistically limited

Charm lifetimes

Charm lifetime measurements at Belle II

- World's most precise measurement of D^0/D^+ lifetime, PRL 127, 211801 (2021)


Belle II	World average
$\tau(D^0) = 410.5 \pm 1.1 \text{ (stat.)} \pm 0.8 \text{ (syst.) fs}$	$410.1 \pm 1.5 \text{ fs}$
$\tau(D^+) = 1030.4 \pm 4.7 \text{ (stat.)} \pm 3.1 \text{ (syst.) fs}$	$1040 \pm 7 \text{ fs}$



$$\tau(D^0) = 410.5 \pm 1.1 \text{ (stat.)} \pm 0.8 \text{ (syst.) fs}$$

($202.4 \pm 3.1 \text{ fs PDG}$)


World's most precise measurement





Dark Sector

Dark Sector Search at Belle II

- Light dark matter search, low background, 3D momentum conservation at Belle II
 - Sensitivity for MeV-GeV scenarios
 - Typical processes
 - $e^+ + e^- \rightarrow \text{SM particle} + \text{mediator}$
 - $B \rightarrow \text{SM particle} + \text{mediator}$





- Belle or BaBar did not search for some of the processes (trigger setting, etc.), Belle II already published 2 results with initial data
 - $e^+e^- \rightarrow \mu^+\mu^-Z'$, $Z' \rightarrow \text{invisible}$ (0.28 fb^{-1}) [PRL 124 \(2020\), 141801](#)
 - $e^+e^- \rightarrow a(\rightarrow \gamma\gamma)\gamma$ (**Axion-Like Particle**) (0.44 fb^{-1}) [PRL 125 \(2020\), 161806](#)



Dark Higgsstrahlung




- U(1)' extension of the standard model
 - Massive **dark photon** (A') as the mediator
 - Spontaneous symmetry breaking introduce a **dark Higgs** (h')
 - Phys. Rev. D 79, 115008 (2009)
 - A' couples to SM only via kinetic mixing (ϵ)
 - α_D dark coupling constant

- Mass hierarchy scenarios
 - $m_{h'} > m_{A'}: h' \rightarrow A'A'^{(*)}$, 4had., $2\ell + 2$ had. (final state: 6 tracks), probed by BaBar (2012), Belle (2015)
 - $m_{h'} < m_{A'}: h'$ “long lived thus invisible” (2 tracks), partly probed by KLOE (2015)




Dark Higgstrahlung results

- No significant deviation from SM background expectation is observed (8.34 fb^{-1})
- Upper limits are set on σ and $\epsilon^2 \alpha_D$:
 - Covered region: $1.65 < M_{A'} < 10.51 \text{ GeV}$ and $M_{h'} < M_{A'}$
 - 90% CL UL on σ from 1.7 to 5 fb @ $4 < M_{A'} < 9 \text{ GeV}$
 - For $M_{A'} < 4 \text{ GeV}$: low sensitivity due to trigger eff.
 - For $M_{A'} > 9 \text{ GeV}$: large dimuon background



Summary and prospects


- Belle II already accumulated 380 fb^{-1} data
- Recent results presented in today's talk
 - Semileptonic B decays
 - First $|V_{ub}|$ and $|V_{cb}|$ measurement
 - B^0 lifetime and mixing frequency: next step $\sin(2\phi_1)$
 - $B^+ \rightarrow \rho^+ \rho^0$ measurement for ϕ_2/α
 - Measurement of ϕ_3/γ : first Belle + Belle II analysis
 - Lifetime of D^0 , D^+ and Λ_c^+ : world most precise
 - Search for Dark sector
- Still other results can not covered in this talk
- More results are expected with competitive intergraded luminosity of BaBar experiment plan to be taken until summer 2022



Backup

B rare decay

$B \rightarrow K\pi$ puzzle



- Assume penguin + tree diagrams are dominant

$$A_{CP}(K^+\pi^-) + \frac{A_{CP}(K^0\pi^+)}{\text{small}} \approx A_{CP}(K^+\pi^0) + \frac{A_{CP}(K^0\pi^0)}{\text{small}}$$

$$A_{CP}^{K^+\pi^0} \sim A_{CP}^{K^+\pi^-}$$

- Current results obtained from the experiment

$$\Delta A_{K\pi} = A_{CP}^{K^+\pi^0} - A_{CP}^{K^+\pi^-} = \mathbf{(12.4 \pm 2.1)\%} \quad \text{PDG}$$

- Another approach (isospin sum rule) to pin down the $B \rightarrow K\pi$ puzzle (less theoretical uncer.)
- QCD color suppression effect cancell out



$$I_{K\pi} \equiv A_{CP}^{K^+\pi^+} + A_{CP}^{K^0\pi^+} \frac{\mathcal{B}_{K^0\pi^+}}{\mathcal{B}_{K^+\pi^-}} \frac{\tau_{B^0}}{\tau_{B^+}} - 2 A_{CP}^{K^+\pi^0} \frac{\mathcal{B}_{K^+\pi^0}}{\mathcal{B}_{K^+\pi^-}} \frac{\tau_{B^0}}{\tau_{B^+}} - A_{CP}^{K^0\pi^0} \frac{\mathcal{B}_{K^0\pi^0}}{\mathcal{B}_{K^+\pi^-}} = 0(?)$$

World average: $I_{K\pi} = (-14 \pm 11)\%$

Neutral final states are crucial !

Measurement of $B \rightarrow K\pi$ decays

- Perform 4D fit (ΔE , Δt , M_{bc} , continuum suppression output)
- Constrain S_{CP} using previous measurements to maximize precision on A_{CP}







	$B \rightarrow K^0\pi^0 (189.8 \text{ fb}^{-1})$
$\text{Br}(10^{-6})$	$11.0 \pm 1.2 \text{ (stat.)} \pm 1.0 \text{ (syst.)}$
PDG(10^{-6})	9.9 ± 0.5
A_{CP}	$-0.41^{+0.30}_{-0.32} \pm 0.09$
$A_{CP}(\text{PDG})$	0.00 ± 0.13

The Belle II Physics Book, PTEP 2019, 123C01

Uncertainty ~4% at Belle II, able to answer $I_{K\pi}$


Dark Sector Search at Belle II

- Belle or BaBar did not search for some of the processes (trigger setting, etc.),
Belle II initial data enable two searches
 - $e^+e^- \rightarrow \mu^+\mu^-Z'$, $Z' \rightarrow \text{invisible}$ (0.28 fb^{-1}) PRL 124 (2020), 141801
 - $e^+e^- \rightarrow a(\rightarrow \gamma\gamma)\gamma$ (**Axion-Like Particle**) (0.44 fb^{-1}) PRL 125 (2020), 161806




Dark Higgstrahlung background and systematics

- **Backgrounds**
 - dominant backgrounds:
 - $\mu^+\mu^-(\gamma)$ (79%)
 - $\tau^+\tau^-(\gamma)$ (18%)
 - $e^-e^+\mu^-\mu^+$ (3%)
 - different contributions in different regions
- **Systematics:**
 - impacting both signal and background: 2.2%-12.7%
 - impacting signal only:
 - differences in M resolution in data/MC (1-5%),
BR theory uncert. 4%




Unitarity Triangle fit extrapolation

The Belle II Physics Book, PTEP 2019, 123C01




ϕ_2 measurement ($B \rightarrow \rho\rho$)



f_L = fraction of longitudinally polarized events

- Compatible with PDG value
- Performance superior to early Belle results



- $\Delta\phi_2 \sim 0.6^\circ$ (current 4.2°) with 50 fb^{-1} data
- $B \rightarrow \pi\pi$, $B \rightarrow \rho\rho$ isospin analysis and $B \rightarrow \rho(\pi\pi)\pi$ Dalitz analysis of 3 body decays
- **LHCb can not measure ϕ_2**

Belle II - LHCb comparison

P. URQUIJO @ Beauty 2020

Belle II
Higher sensitivity to decays with photons and neutrinos (e.g. $B \rightarrow K\bar{v}v, \mu\nu$), inclusive decays, time dependent CPV in B_d, τ physics.
LHCb
Higher production rates for ultra rare $B, D, & K$ decays, access to all b -hadron flavours (e.g. Λ_b), high boost for fast B_s oscillations.
<i>Overlap in various key areas to verify discoveries.</i>
Upgrades
<i>Most key channels will be stats. limited (not theory or syst.).</i> LHCb scheduled major upgrades during LS3 and LS4. Belle II formulating a 250 ab^{-1} upgrade program post 2028.


Observable	Current Belle/ Babar	2019 LHCb	Belle II (5 ab ⁻¹)	Belle II (50 ab ⁻¹)	LHCb (23 fb ⁻¹)	Belle II Upgrade (250 ab ⁻¹)	LHCb upgrade II (300 fb ⁻¹)
<u>CKM precision, new physics in CP Violation</u>							
$\sin 2\beta/\phi_1 (B \rightarrow J/\psi K_S)$	0.03	0.04	0.012	0.005	0.011	0.002	0.003
γ/ϕ_3	13°	5.4°	4.7°	1.5°	1.5°	0.4°	0.4°
α/ϕ_2	4°	—	2	0.6°	—	0.3°	—
$ V_{ub} $ (Belle) or $ V_{ub} / V_{cb} $ (LHCb)	4.5%	6%	2%	1%	3%	<1%	1%
Φ_s	—	49 mrad	—	—	14 mrad	—	4 mrad
$S_{CP}(B \rightarrow \eta' K_S, \text{gluonic penguin})$	0.08	○	0.03	0.015	○	0.007	○
$A_{CP}(B \rightarrow K_S \pi^0)$	0.15	—	0.07	0.04	—	0.02	—
<u>New physics in radiative & EW Penguins, LFUV</u>							
$S_{CP}(B_d \rightarrow K^* \gamma)$	0.32	○	0.11	0.035	○	0.015	○
$R(B \rightarrow K^* l^+ l^-) (1 < q^2 < 6 \text{ GeV}^2/c^2)$	0.24	0.1	0.09	0.03	0.03	0.01	0.01
$R(B \rightarrow D^* \tau \nu)$	6%	10%	3%	1.5%	3%	<1%	1%
$Br(B \rightarrow \tau \nu), Br(B \rightarrow K^* \bar{v}v)$	24%, —	—	9%, 25%	4%, 9%	—	1.7%, 4%	—
$Br(B_d \rightarrow \mu \mu)$	—	90%	—	—	34%	—	10%
<u>Charm and τ</u>							
$\Delta A_{CP}(K\bar{K} - \pi\pi)$	—	8.5×10^{-4}	—	5.4×10^{-4}	1.7×10^{-4}	2×10^{-4}	0.3×10^{-4}
$A_{CP}(D \rightarrow \pi^+ \pi^0)$	1.2%	—	0.5%	0.2%	—	0.1%	—
$Br(\tau \rightarrow e \gamma)$	$< 120 \times 10^{-9}$	—	$< 40 \times 10^{-9}$	$< 12 \times 10^{-9}$	—	$< 5 \times 10^{-9}$	—
$Br(\tau \rightarrow \mu \mu \mu)$	$< 21 \times 10^{-9}$	$< 46 \times 10^{-9}$	$< 3 \times 10^{-9}$	$< 3 \times 10^{-9}$	$< 16 \times 10^{-9}$	$< 0.3 \times 10^{-9}$	$< 5 \times 10^{-9}$

Results on other D & τ
modes expected

Prospects of $|V_{ub}|$ and $|V_{cb}|$

The Belle II Physics Book, PTEP 2019, 123C01

Side	Observable	Dominant uncertainties
$ V_{td} $	Δm_d : $B\bar{B}$ mixing frequency	Lattice QCD ($ V_{td} $ now is mainly limited by LQCD)
$ V_{cb} $	$Br(b \rightarrow c/\ell\nu)$	Exclusive: Lattice QCD Inclusive: experiment vs. phenomenology
$ V_{ub} $	$Br(b \rightarrow u/\ell\nu)$	



Observables	Belle (2017)	Belle II	
		5 ab^{-1}	50 ab^{-1}
$ V_{cb} $ incl.	$42.2 \cdot 10^{-3} \cdot (1 \pm 1.8\%)$	1.2%	—
$ V_{cb} $ excl.	$39.0 \cdot 10^{-3} \cdot (1 \pm 3.0\%_{\text{ex.}} \pm 1.4\%_{\text{th.}})$	1.8%	1.4%
$ V_{ub} $ incl.	$4.47 \cdot 10^{-3} \cdot (1 \pm 6.0\%_{\text{ex.}} \pm 2.5\%_{\text{th.}})$	3.4%	3.0%
$ V_{ub} $ excl. (WA)	$3.65 \cdot 10^{-3} \cdot (1 \pm 2.5\%_{\text{ex.}} \pm 3.0\%_{\text{th.}})$	2.4%	1.2%
$\mathcal{B}(B \rightarrow \tau\nu) [10^{-6}]$	$91 \cdot (1 \pm 24\%)$	9%	4%
$\mathcal{B}(B \rightarrow \mu\nu) [10^{-6}]$	< 1.7	20%	7%
$R(B \rightarrow D\tau\nu)$ (Had. tag)	$0.374 \cdot (1 \pm 16.5\%)$	6%	3%
$R(B \rightarrow D^*\tau\nu)$ (Had. tag)	$0.296 \cdot (1 \pm 7.4\%)$	3%	2%

Belle II - LHCb comparison

