

The Belle II Experiment: Status and Prospects

K. Lalwani

(on behalf of the Belle II Collaboration)

Malaviya National Institute of Technology Jaipur, Jaipur, India
kavita.phy@mnit.ac.in

August 11, 2021

1 Abstract

The Belle II experiment at the SuperKEKB energy asymmetric e^+e^- collider is a substantial upgrade of the B factory facility at the Japanese KEK laboratory. The design luminosity of the machine is $6 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ and the Belle II experiment aims to record 50 ab^{-1} of data, a factor of 50 more than its predecessor. With this data set, Belle II will be able to measure the Cabibbo-Kobayashi-Maskawa (CKM) matrix, the matrix elements and their phases, with unprecedented precision and explore flavor physics with B , charmed mesons, and τ leptons. Belle II has also a unique capability to search for low mass dark matter and low mass mediators. In this paper, we will review the status of the Belle II detector, SuperKEKB accelerator and the prospects for physics at Belle II.

13 1 Introduction

Heavy flavour physics plays a key role in understanding the Standard Model (SM) and its mechanism. The first generation of B factories [1], KEKB, PEP-II and their related experiments Belle and BaBar successfully operated for 10 years and achieved substantial physics results. Both experiments provided significant contributions to B physics in finding the first evidence of CP violation outside the kaon system [2] and the experimental confirmation of the Cabibbo-Kobayashi-Maskawa (CKM) mechanism [3]. There are still several SM predictions, which need to be verified, and the investigation of New Physics (NP) processes is extremely important. Therefore, a second generation B factory with a low-background environment and large data samples of B , D , and τ is needed, which will have exclusive advantages as compared to the hadronic machines. The Belle II experiment [4] at SuperKEKB [5] is the successor to the previous Belle experiment at KEKB. The design luminosity of SuperKEKB is $6 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ and the Belle II experiment aims to record 50 ab^{-1} of data, which is a factor of 50 more than its predecessor. With this huge data set, Belle II is expected to extend the search for NP in the flavour sector at the precision frontier using a complementary approach with respect to LHC experiments. This paper reviews the status of the Belle II experiment and SuperKEKB. The latest results on B physics, charm physics and τ physics at Belle II are also discussed.

31 2 SuperKEKB Accelerator

The SuperKEKB accelerator machine is situated at the High Energy Accelerator Research Organization (KEK) in Tsukuba, Japan. The design luminosity of the SuperKEKB ac-

34 celerator is $6 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$, which is 40 times greater than that of the KEKB. In order
 35 to achieve this high luminosity, a nano beam scheme [6] is introduced by SuperKEKB,
 36 where luminosity is greatly increased by increasing the beam current (by a factor ~ 2)
 37 and reducing the vertical beta function at the IP (by a factor ~ 20). However with the
 38 increase in luminosity, the beam related background also increases, which will be handled
 39 with the improved Belle II detector. The SuperKEKB accelerator (figure 1 (left)) reached
 to the world record peak luminosity of $2.40 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ for an e^+e^- collider.




Figure 1: SuperKEKB accelerator (left) and Belle II detector (right).

40

41 3 Belle II Detector

42 Due to the high luminosity of SuperKEKB, the Belle II detector will be operated in a
 43 harsher radiation environment compared to Belle. In order to cope with this high back-
 44 ground, almost all Belle II sub-detectors have been substantially upgraded. A new vertex
 45 detector consisting of a Pixel vertex detector and four layers of fast Silicon vertex detec-
 46 tor is introduced, which provides the improved vertex resolution by a factor of two as
 47 compared to Belle. Further, we have a new Central Drift Chamber (CDC), which is the
 48 main tracking detector and it provides the better charge track reconstruction and dE/dx
 49 measurement. It is built with smaller cells than Belle's to operate with higher event rates.
 50 Outside the CDC, we have a particle identification (PID) system consisting of a Time-
 51 of-Propagation Counter in the barrel region and the Aerogel Ring-Imaging Cherenkov
 52 detector in the forward-end-cap region, which are mainly used to distinguish pions from
 53 kaons with a fake rate lower than in Belle. After the PID system, we have an electromag-
 54 netic calorimeter, which is substantially the same as used in Belle detector, with a faster
 55 read-out electronics. A K_L meson and μ detector has been improved by substituting all
 56 the Resistive Plate Chamber layers with scintillators in the end-caps region and the first
 57 two layers in the barrel region. The upgraded Belle II detector (figure 1 (right)) is ex-
 58 pected to provide improved impact parameter resolution, increased K_s efficiency, a better
 59 K/π separation and good π^0 reconstruction. Belle II has recorded data corresponding to
 60 an integrated luminosity of 213.49 fb^{-1} .

61 4 Performance of the Belle II Detector

62 The performance of the Belle II detector is validated using various control samples. The
 63 performance of the charged kaon and pion identification is studied using data correspond-

ing to an integrated luminosity of $37fb^{-1}$. The results of kaon efficiency and pion mis-ID rates for different PID criteria using the decay $D^{*+} \rightarrow D^0[K^-\pi^+]\pi^+$ are shown in figure 2 (left). This study is performed in several bins of laboratory frame momentum and polar angle [7]. The tracking efficiency and fake rate are measured using $e^+e^- \rightarrow \tau^+\tau^-$ events in e^+e^- collision data collected in 2019 at Belle II, where one tau lepton decays leptonically ($\tau \rightarrow \ell^\pm\nu_\ell\bar{\nu}_\tau, \ell = e, \mu$), while the other decays hadronically into three charged pions ($\tau \rightarrow 3\pi^\pm\nu_\tau + n\pi^0$) [8] as shown in figure 2 (right). Further, reconstruction performance of neutral particles at Belle II is demonstrated by analysing the two photon events coming from π^0 and η [9].




Figure 2: K -identification efficiencies and π -misidentification rates for different PID criteria using the decay $D^{*+} \rightarrow D^0[K^-\pi^+]\pi^+$ (left), measured tracking efficiency times detector acceptance ($\epsilon \times A$) and calibrated data-MC discrepancy (δ^*) for the combined channels as a function of the 1-prong track p_T (right).

72

73 5 Physics Programme at Belle II

74 The Belle II experiment aims to investigate heavy flavour physics with high precision as
 75 a B factory. Physics programme at Belle II covers wide range of physics, which includes
 76 B , D and τ leptons along with dark sector searches. In this paper, important highlights
 77 on limited physics studies such as measurement of the CKM angles, time integrated CP
 78 asymmetry using charmless B decays, D^0 lifetime along with τ -mass measurement will be
 79 discussed.

80 5.1 Measurement of the CKM Angles

81 Due to good flavor tagging efficiency at Belle II, it provides an opportunity to study
 82 CP violation by measuring the CKM angles through various B decays; discrimination
 83 of signal from background utilizes two important variables ΔE (beam-energy difference)
 84 and M_{bc} (beam-constrained mass). The decay $B^0 \rightarrow J/\psi K_L^0$ provides an independent
 85 measurement of CKM angle $\sin(2\phi_1)$, where J/ψ is reconstructed from e^+e^- and $\mu^+\mu^-$,
 86 and K_L^0 is reconstructed as a hadronic neutral cluster in KLM. Figure 3 (top: left) shows
 87 ΔE distribution for $B^0 \rightarrow J/\psi K_L^0$ with data corresponding to an integrated luminosity
 88 of $62.8fb^{-1}$. Figure 3 (top: right) shows M_{bc} distribution for $B^0 \rightarrow J/\psi K_S^0$ with data
 89 corresponding to an integrated luminosity of $34.6fb^{-1}$. Further, ΔE distribution for $B^0 \rightarrow$
 90 $\pi^0\pi^0$ is shown in figure 3 (bottom: left), which is difficult to reconstruct, as it has four
 91 photons in final state. This decay is important to measure the CKM angle (ϕ_2). Figure 3

92 (bottom: right) shows the ΔE distribution for $B^0 \rightarrow D^0 h^-$, where h is either a kaon or a
 93 pion. This study is aimed to measure the CKM angle (ϕ_3) with higher precision [10].




Figure 3: ΔE distribution for $B^0 \rightarrow J/\psi K_L^0$ (top: left), M_{bc} distribution for $B^0 \rightarrow J/\psi K_s^0$ (top: right), ΔE distribution for $B^0 \rightarrow \pi^0 \pi^0$ (bottom: left), and ΔE distribution for $B^0 \rightarrow D^0 h^-$, where h is either a kaon or a pion (bottom: right).

94 5.2 $B \rightarrow K\pi$ decays

95 The $K\pi$ isospin sum rule [11] offers a stringent null test of the SM, and is expressed
 96 in terms of direct CP asymmetries and branching fractions of the four $B \rightarrow K\pi$ decay
 97 modes. We observed 45_{-8}^{+9} signal events from the fitting of ΔE and M_{bc} distributions of
 98 $B^0 \rightarrow K^0 \pi^0$, which is translated to $\mathcal{B}(B^0 \rightarrow K^0 \pi^0) = (8.5_{-1.6}^{+1.7} \pm 1.2) \times 10^{-6}$ [12]. As this
 99 decay is a CP eigen-state, we use the output of the flavor tagger to determine the time
 100 integrated CP asymmetry $[-0.40_{-0.44}^{+0.46} \pm 0.04]$ [12].

101 5.3 Measurement of D^0 life time

102 The lifetime measurement of D^0 meson is performed with data corresponding to an inte-
 103 grated luminosity of $9.6 fb^{-1}$ using the three decays modes, namely, $D^0 \rightarrow K^- \pi^+$, $D^0 \rightarrow$
 104 $K^- \pi^+ \pi^0$ and $D^0 \rightarrow K^- \pi^+ \pi^- \pi^+$ coming from $D^{*+} \rightarrow D^0 \pi^+$ [13]. The D^0 lifetime is
 105 measured by performing a two-dimensional unbinned ML fit to distributions of proper
 106 time and its uncertainty (figure 4 (left)). The average lifetime of the D^0 meson is mea-
 107 sured to be $(412.3 \pm 2.0) fs$ (figure 4 (right)). With $72 fb^{-1}$ of Belle II data, the lifetime
 108 measurement of the D^0 meson is expected to be competitive with the world-averages.

109 5.4 Preliminary analysis of charm meson decays

110 Due to the large data sample of charm mesons produced at Belle II, it is a good opportunity
 111 to investigate the CP violation (CPV) in the charm sector as well. In particular, the
 112 time-integrated Dalitz plot analysis of $D^{*+} \rightarrow D^0[\rightarrow \pi^+ \pi^- \pi^0] \pi^+$ mode could be used
 113 to search for CPV. A signal yield of $305 \pm 15(stat.)$ is extracted using the distribution of
 114 $\Delta M = m(D^*) - m(D^0)$ with data corresponding to an integrated luminosity of $72 fb^{-1}$ [14]




Figure 4: Proper-time distribution of the D^* tagged candidates in the $D^0 \rightarrow K^-\pi^+$ channel (left), comparison of the D^0 life time at Belle II with the world average values (right).

115 as shown in figure 5 (left). In addition, rediscovery of Singly Cabibbo Suppressed (SCS)
 116 decay $D^0 \rightarrow K_s K_s$ is also carried out at Belle II [15]. Further, the ratios of wrong side
 117 to right side (WS to RS) yield of three decay modes ($D^0 \rightarrow K^+\pi^-$, $D^0 \rightarrow K^+\pi^-\pi^0$ and
 118 $D^0 \rightarrow K^+\pi^-\pi^+\pi^-$) are also measured and results are in agreement with PDG values [16]
 as shown in figure 5 (right).




Figure 5: ΔM distribution of $D^{*+} \rightarrow D^0[\rightarrow \pi^+\pi^-\pi^0]\pi^+$ (left) and ratio of WS to RS yield (right).

119

120 5.5 Tau mass measurement

121 The measurement of mass of τ lepton is carried out at Belle II with data corresponding
 122 to an integrated luminosity of 8.8 fb $^{-1}$ [17]. The tau mass is measured to be $1777.28 \pm$
 123 $0.75(stat.) \pm 0.33(syst.)$ MeV. The precision of this measurement is limited by the size
 124 of the data that was used, but the systematic uncertainty is comparable to that at Belle.
 125 With further data provided by the Belle II experiment, the statistical uncertainty will
 126 further decrease.

127 6 Summary

128 Belle II has been running continuously and collecting data despite the Covid-19 pandemic.
 129 Its aim is to record an integrated luminosity of 50 ab $^{-1}$. This upcoming large and clean
 130 data samples of B and D mesons (and τ leptons) will allow Belle II to search for NP and
 131 improve the measurements of various SM parameters. The results reported in this paper

132 are based on early Belle II data and show the Belle II's performance is as expected.

133 References

- 134 [1] A. J. Bevan *et al.*, *The Physics of the B Factories*, Eur. Phys. J. C **74**, 3026 (2014).
- 135 [2] K. Abe *et al.*, *Observation of large CP violation in the neutral B meson system*, Phys.
136 Rev. Lett. **87**(9), 091802 (2001).
- 137 [3] M. Kobayashi *et al.*, *CP Violation in the renormalizable theory of weak interaction*,
138 Prog. Theor. Phys. **49**, 652 (1973).
- 139 [4] E. Kou *et al.*, *The Belle II physics book*, Prog. Theor. Exp. Phys. **2019**(12), 123C01
140 (2019).
- 141 [5] K. Akai *et al.*, *SuperKEKB collider*, Nucl. Instrum. Meth. A **907**, 188 (2018).
- 142 [6] P. Raimondi *et al.*, *Proceedings of 11th European Conference*, EPAC 2008, Genoa,
143 Italy **2620**, 652 (2008).
- 144 [7] Belle II collaboration, *Kaon and Pion Identification Performance in Phase III data*,
145 BELLE2-NOTE-PL-2020-024.
- 146 [8] Belle II collaboration, *Measurement of the tracking efficiency and fake rate with*
147 *$e^+e^- \rightarrow \tau^+\tau^-$ events*, BELLE2-NOTE-PL-2020-014.
- 148 [9] Belle II collaboration, *plot of $\pi^0 \rightarrow \gamma\gamma$ and $\eta \rightarrow \gamma\gamma$ in Early Phase 3 Belle II data*,
149 BELLE2-NOTE-PL-2019-019.
- 150 [10] Belle II collaboration, *Study of $B \rightarrow D^- h$ decays using 62.8 fb^{-1} of Belle II data*,
151 BELLE2-NOTE-PL-2019-028.
- 152 [11] M. Gronau *et al.*, *Combining CP asymmetries in $B \rightarrow K\pi$ decays*, Phys. Rev. D **59**,
153 113002 (1999).
- 154 [12] Belle II collaboration, *First search for direct CP-violating asymmetry in $B^0 \rightarrow K^0\pi^0$*
155 *decays at Belle II*, BELLE2-CONF-PH-2021-001.
- 156 [13] Belle II collaboration, *D^0 Lifetime Plots with 2019 Data*, BELLE2-NOTE-PL-2020-
157 008.
- 158 [14] Belle II collaboration, *Preliminary analysis of $D^{*+} \rightarrow D^0(\pi^+\pi^-\pi^0)\pi^+$* , BELLE2-
159 NOTE-PL-2021-003.
- 160 [15] Belle II collaboration, *Rediscovery of the $D^0 \rightarrow K_s K_s$ decay with early Belle II data*,
161 BELLE2-NOTE-PL-2020-020.
- 162 [16] Belle II collaboration, *Rediscovery of "wrong-sign" D^0 decays to hadronic final states*
163 *with early Belle II data*, BELLE2-NOTE-PL-2020-021.
- 164 [17] Belle II collaboration, *τ lepton mass measurement at Belle II*, BELLE2-CONF-PH-
165 2020-010.