

## Recent quarkonium results at Belle II

---

Dmytro Meleshko (*on behalf of Belle II collab.*)

October 11, 2022




Justus-Liebig-Universität, Giessen, Germany

Excited QCD, Sicily, Italy.




# Quarkonium(-like) mess

- Below  $D\bar{D}/B\bar{B}$  thresholds  $c\bar{c}$  and  $b\bar{b}$  match QCD;
- Many exotic states observed in the past decade are hard to fit these spectra.



# Belle II experiment overview




## SuperKEKB:

- Asymmetric  $e^+e^-$  collider at KEK (Tsukuba, Japan);
- Energy adjustment: 3.5/8.0 GeV (Belle)  $\rightarrow$  7.0/4.0 GeV (Belle II);
- "Nano-beams"  $\times$  current increase ( $\times 2$ ) =  $\times 40$  inst. luminosity increase;


## Belle II detector upgrade:

- Higher background:
  - Radiation damage;
  - Detector readout;
- Higher event rate:
  - Trigger, DAQ, computing;
- Boost change:
  - Vertexing improvement;




# Belle II timeline

- 2016: "Phase 1":
  - Beam commissioning;
- 2017: Detector roll-in;
- 2018: "Phase 2":
  - Background study w. partial detector;
  - First collisions/data (28.04.2018);
- 2019: "Phase 3":
  - Nominal start of operations;
  - 2021: inst. lumin. record:  $> 4.7 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$ ;
  - 2021: Non  $\Upsilon(4S)$  energy scan;
- 2022-2023: "Long Shutdown 1":
  - Detector/accelerator upgrades;
- 2023-2027: Resume operations, target:  $5 \text{ab}^{-1}$ ;
- 2027+: "Long Shutdown 2" upgrade (?), continue up to  $50 \text{ab}^{-1}$ .



# Belle II quarkonium potential

- Many flavor physics contributions, particularly in hadron spectroscopy;
- Advantages:
  - “Clean” environment;
  - Full event reconstruction, decays with neutral/soft particles;
  - Nominal  $\sqrt{s} = 10.58 \text{ GeV} = m(\Upsilon(4S))$ , potential to reach 11 GeV;
  - Decay with neutrals ( $\gamma, \pi^0, K_l, \nu$ ) in final state;
  - Multiple production mechanisms;
  - Large statistics;
  - Complementary to LHC.






## Charmonium(-like) studies at Belle II

---

# Charmonium(-like) studies at Belle II: $B$ -decays

## $B$ -decays

- $B \rightarrow KX_{c\bar{c}}$ : CKM favored process  $\rightarrow$  large branching fractions ( $10^{-3} \sim 10^{-4}$ );
- Absolute  $\mathcal{B}_f(B \rightarrow X(3872, 3915)K)$ ;
- $X(3872)$  width measurement with  $D^0 \bar{D}^0 \pi^0$ ;
- Confirm  $Z_c$  states and search for neutral partners.







First observation of  $X(3872)$  at Belle II:  $14.4 \pm 4.6$  events ( $4.6\sigma$ )

\*  $\epsilon(\psi(2S)) = 20\%$   
(w.r.t. Belle)

| State     | Production and decay   | N               |
|-----------|--|-----------------|
| $X(3872)$ | $B \rightarrow KX(3872)$ , $X(3872) \rightarrow J/\psi \pi^+ \pi^-$  | $\approx 14400$ |
| $Y(4260)$ | ISR, $Y(4260) \rightarrow J/\psi \pi^+ \pi^-$                        | $\approx 29600$ |
| $Z(4430)$ | $B \rightarrow K^\mp Z(4430)$ , $Z(4430) \rightarrow J/\psi \pi^\pm$ | $\approx 10200$ |

# Charmonium(-like) studies at Belle II: $\Gamma(X(3872))$

- Features:
  - Study of  $X(3872)$  in  $D^0\bar{D}^0\pi^0$  is a new approach;
  - Extremely small Q value gives an advantage;
- Possibilities:
  - $(684 \pm 8)$  keV mass resolution (vs.  $1.93 \pm 0.04$  for  $J\psi\pi\pi$ );
  - Push  $\Gamma^{UL}(X(3872))$  down to  $\approx 280$  keV (Note: Flatté fit  $220^{+70+11-}_{-60-130}$ );
  - Allows to decrease systematic uncertainty down to 110 keV;
  - possibility to combine  $D^0\bar{D}^0\pi^0$  and  $J/\psi\pi\pi$ .
- Another idea: search for exotics at  $D^*\bar{D}^*$  threshold.





| Mode                | Q value (MeV)     |
|---------------------|-------------------|
| $J/\psi\pi^+\pi^-$  | $495.65 \pm 0.17$ |
| $D^0\bar{D}^0\pi^0$ | $7.05 \pm 0.18$   |
| $D^0\bar{D}^0*$     | $0.01 \pm 0.18$   |



# Charmonium(-like) studies at Belle II: other processes

## Initial-state radiation (ISR):

- Continuous mass range  $>4.7 \text{ GeV}/c^2$ ;
- $Y \rightarrow c\text{-baryon pairs } (\Lambda_c^+, \Sigma_c^-, \Sigma_c^+ \Sigma_c^-)$ ,  $cs\text{-meson pairs } (D_s D_{s2}(2573), D_s D_{s0}^*(2317))$ ;
- Search for  $Z_{cs}$  states decaying into  $K^\pm J/\psi$ ,  $D_s^- D^{*0} + \text{c.c.}$ ;
- $Y(4260)$  rediscovery (expected 60 events per  $100 \text{ fb}^{-1}$ ) + line shape;




\* clear observation of very clean ISR  $\psi(2S)$  signals

## Two-photon process:


- $J^{PC}$  of  $X(3915) \rightarrow \omega J/\psi$ ;
- Confirm  $X(4350) \rightarrow \phi J/\psi$ ;

## Double charmonium:


- $e^+ e^- \rightarrow (c\bar{c})_{J=1} (c\bar{c})_{J=0}$  production rule;
- $J^{PC}$  of  $X(3940)$ .



Initial state radiation



Double  $c\bar{c}$





$\gamma\gamma$  collisions

## **Bottomonium(-like) studies at Belle II**

---


# Bottomonium: life outside $\Upsilon(4S)$

- B-factories' advantages for bottomonium research:
  - Able to adjust  $E_{CM}$  in 9.4-11.2 GeV region;
  - Pole-position for  $\Upsilon$ ,  $Y_b$  and  $Z_b$  states.
- B-factories legacy in bottomonium field:
  - BaBar  $\Upsilon(3S)$ : discovery of  $\eta_b(1S)$   
arXiv:0809.1672v1 [hep-ex] 9 Sep 2008;
  - Belle  $\Upsilon(5S)$ : dicoverly of  $h_b(1P, 2P)$ ,  $\eta_b(2S)$ ,  $Z_b(10610, 10650)^\pm$   
PRL108,032001(2012), arXiv:1205.6351, arXiv:1105.5492;
  - Belle:  $Y_b(10753)$  discovery  
JHEP 06 (2019) 220;





- Belle II data acquisition target:
  - 100 fb<sup>-1</sup> at  $\Upsilon(6S)$ ;
  - 1 ab<sup>-1</sup> at  $\Upsilon(5S)$ .
- Principal ROI:
  - $\Upsilon(6S)$  vs.  $\Upsilon(5S)$  in  $\pi\pi\Upsilon$  and  $\pi\pi\eta$  decays: is  $\Upsilon(5S)$  a  $b\bar{b}$  state?
  - Deeper study of 10.750 GeV/c<sup>2</sup> vicinity;
  - $c\bar{c}$  vs.  $b\bar{b}$  spectrum discrepancy;

# Bottomonium: Belle II early physics results






Prelude to energy scan analysis

- ISR processes:
  - $\gamma_{ISR} \Upsilon(2S) \rightarrow \pi^+\pi^-\Upsilon(1S)$ ;
  - $\gamma_{ISR} \Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(1S, 2S)$ ;
- Direct transitions:  $\Upsilon(4S) \rightarrow \pi^+\pi^-\Upsilon(1S, 2S)$ ;
- First-look results:
  - Improvement w.r.t. Belle;
  - The 3S → 2S transition is seen;
  - $\Upsilon(4S) \rightarrow \pi^+\pi^-\Upsilon(nS)$  Datitz analysis is ongoing;






# Belle II Energy Scan (Nov 2021)

- Considerations:
  - Early physics at Belle II: fruitful field.
  - Major constraint: limited luminosity:  $O(15\text{fb}^{-1})$ .
  - Prospects: cover  $\Upsilon(6S)$  region and beyond after detector upgrade.
- Status:
  - Energy scan operation was successful: x4 higher statistic w.r.t. Belle is accumulated.




# Belle II potential: 10.75 GeV




- Belle: 7 energy scan datasets ( $1 \text{ fb}^{-1}$  each) below  $\Upsilon(5S)$ .
- New structure is observed in  $\Upsilon(nS)\pi^+\pi^-$  system ( $n=1,2,3$ ;  $\Upsilon(nS) \rightarrow e^+e^-/\mu^+\mu^-$ ).
- Not seen in  $B^{(*)}\bar{B}^{(*)}$  cross section (more data?).

| Name              | Properties [MeV]   |
|-------------------|--|
| $\Upsilon(10860)$ | $M = 10885.3 \pm 1.5^{+2.2}_{-0.9}$<br>$\Gamma = 36.6^{+4.5+0.5}_{-3.9-1.1}$   |
| $\Upsilon(11020)$ | $M = 11020.3^{+4.0+1.0}_{-4.5-1.3}$<br>$\Gamma = 23.8^{+8.0+0.7}_{-6.8-1.8}$   |
| New structure     | $M = 10752.7 \pm 5.9^{+0.7}_{-1.1}$<br>$\Gamma = 35.5^{+17.6+3.9}_{-11.3-3.3}$ |



# Why is $Y_b(10753)$ important?

- First observation (Belle, 2019) in  $\sigma(e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-)$  vs.  $E$  ( $\sqrt{s} = 10.6 - 11.2$  GeV).  
R. Mizuk et al., J. High Energy Phys. 10, 220 (2019).
- Unclear nature
  - Pure  $\Upsilon(3D)$  interpretation contradicts with theory.  
Godfrey, Moats, Phys. Rev. D 92 (2015) no.5, 054034.
  - Molecule? Too far from the threshold.



- Tetraquark?
- Hybrid?
- ...
- Fast lane to today's aspects of XYZ puzzle in charmonium.

## Conventional $b\bar{b}$ interpretation:

Bai, Li, Huang, Liu, Matsuki, Phys. Rev. D 105, 074007 (2022).  
Li, Bai, Huang, Liu, Phys. Rev. D 104, 034036 (2021).  
Li, Liu, Liu, Gui, Zhong, Eur. Phys. J. C 80, 59 (2020).  
Chen, Zhang, He, Phys. Rev. D 101, 014020 (2020).  
Giron and Lebed, Phys. Rev. D 102, 014036 (2020).  
Kher, Chaturvedi, Devlani, Rai, Eur. Phys. J. Plus 137, 357 (2022).  
Li, Bai, Liu, arXiv:2205.04049.  
Liang, Ikeno, Oset, Phys. Lett. B 803, 135340 (2020).  
Hüsken, Mitchell, Swanson, arXiv:2204.11915.  
Beveren, Rupp, Prog. Part. Nucl. Phys. 117, 103845 (2021).

## Tetraquark interpretation:


Ali, Maiani, Parkhomenko, Wang, Phys. Lett. B 802, 135217 (2020).  
Bicudo, Cardoso, Müller, Wagner, Phys. Rev. D 103, 074507 (2021).  
Bicudo, Cardoso, Müller, Wagner, arXiv:2205.11475.  
Wang, Chin. Phys. C 43, 123102 (2019).

## Hybrid interpretation:


Castella, Passemar, Phys. Rev. D 104, 034019 (2021).  
Brambilla, Eidelman, Hanhart, Nefediev, Shen, Thomas, Vairo, Yuan, Phys. Rept. 873, 1 (2020).

# Motivation to start a new analysis over Belle II scan data


- Theoretical background:
  - $B(Y(10753)) \rightarrow \omega\chi_{bJ}$   $10^{-3}$  for  $4S - 3D$   $b\bar{b}$  mixture.  
Li, Bai, Huang, Liu, Phys. Rev. D 104, 034036 (2021)
- $X_b$ :  $X(3872)$  counterpart candidate?  
Choi et al., Phys. Rev. Lett. 91, 262001 (2003)
  - The  $\pi\pi J/\psi$  and  $\pi\pi\Upsilon$  cross section spectrum similarity hints at similar nature.
  - $Y(4220)$  was observed by BESIII in  $\gamma X(3872)$  and  $\omega\chi_{c0}$  decays.
  - Should we expect  $Y(10753)$  to decay into  $\gamma X(3872)$  and  $\omega\chi_{c0}$ ? Worth checking!



arXiv:1611.01317v2, 20 Jan 2017



PhysRevLett.122.232002



PhysRevD.99.091103



## Observation of $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$ and search for $X_b \rightarrow \omega\Upsilon(1S)$ at $\sqrt{s}$ near 10.75 GeV

I. Adachi, L. Aggarwal, H. Ahmed, H. Aihara, N. Akopov, A. Aloisio, N. Anh Ky, T. Aushev, V. Aushev, H. Bae, P. Bambade, Sw. Banerjee, J. Baudot, M. Bauer, A. Beaubien, J. Becker, P. K. Behera, J. V. Bennett, E. Bernieri, F. U. Bernlochner, V. Bertacchi, M. Bertemes, E. Bertholet, M. Bessner, S. Bettarini, B. Bhuyan, F. Bianchi, T. Bilka, D. Biswas, D. Bodrov, A. Bolz, A. Bondar, J. Borah, A. Bozek, M. Bračko, P. Branchini, T. E. Browder, A. Budano, S. Bussino, M. Campajola, L. Cao, G. Casarosa, M.-C. Chang, P. Cheema, V. Chekelian, Y. Q. Chen, K. Chilikin, K. Chirapatpimol, H.-E. Cho, K. Cho, S.-J. Cho, S.-K. Choi, S. Choudhury, D. Cinabro, L. Corona, S. Cunliffe, S. Das, F. Dattola, E. De La Cruz-Burelo, S. A. De La Motte, G. De Nardo, M. De Nuccio, G. De Pietro, R. de Sangro, M. Destefanis, S. Dey, A. De Yta-Hernandez, R. Dhamija, A. Di Canto, F. Di Capua, Z. Doležal, I. Domínguez Jiménez, T. V. Dong, M. Dorigo, K. Dort, S. Dreyer, S. Dubey, G. Dujany, M. Eliachevitch,



...  
(The Belle II Collaboration)

We study the processes  $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$  ( $J = 0, 1, \text{ or } 2$ ) using samples at center-of-mass energies  $\sqrt{s} = 10.701, 10.745, \text{ and } 10.805$  GeV, corresponding to 1.6, 9.8, and 4.7 fb<sup>-1</sup> of integrated luminosity, respectively. These data were collected with the Belle II detector during a special run of the SuperKEKB collider above the  $\Upsilon(4S)$  resonance. We report the first observation of  $\omega\chi_{bJ}(1P)$  signals at  $\sqrt{s} = 10.745$  GeV. By combining Belle II data with Belle results at  $\sqrt{s} = 10.867$  GeV, we find energy dependencies of the Born cross sections for  $e^+e^- \rightarrow \omega\chi_{b1,b2}(1P)$  to be consistent with the shape of the  $\Upsilon(10753)$  state. Including data at  $\sqrt{s} = 10.653$  GeV, we also search for the bottomonium equivalent of the  $X(3872)$  state decaying into  $\omega\Upsilon(1S)$ . No significant signal is observed for masses between 10.45 and 10.65 GeV/ $c^2$ .

arXiv:2208.13189v1 [hep-ex] 28 Aug 2022

# Analysis Overview

- $e^+e^- \rightarrow \omega\chi_{cJ} (\sigma_b)$ 
  - Born cross section measurement ( $\sigma_b$ ) at  $\sqrt{s} = 10.701, 10.745, 10.805$  GeV.
  - Combining Belle results at  $\sqrt{s} = 10.867$  GeV to study  $\sigma_b$  energy dependence.  
[arXiv:1408.0504v2 \[hep-ex\]](https://arxiv.org/abs/1408.0504v2) 11 Sep 2014
- $e^+e^- \rightarrow \gamma X_b$ 
  - Search for  $X_b$  signal at  $\sqrt{s} = 10.653, 10.701, 10.745, 10.805$  GeV.
- Strategy
  - EvtGen: PHOKHARA and PHSP.
  - Geant4.
  - Offline analysis within Belle II framework.
- Events selection
  - Standard POCA and PID requirements are set charged tracks selection (90%+ eff.).
  - Bremsstrahlung and FSR suppression +  $E(\gamma) > 50$  MeV.
  - Accurate  $\pi^0$  combination.
  - $\chi^2$ -based BCS is applied.



# Observation of $e^+e^- \rightarrow \omega\chi_{bJ}$

- 2D UML fit of  $M(\gamma Y(1S))$  vs.  $M(\pi^+\pi^-\pi^0)$  distribution.
- Model: signal (CB for  $\chi_{bJ}$ , Voigt for  $\omega$ ) + peaking bkg. (same) + comb. bkg.

| Channel                              | $\sqrt{s}$ (GeV) | $N^{sig}$              | $\sigma_B$ (pb)             |
|--------------------------------------|------------------|------------------------|-----------------------------|
| $e^+e^- \rightarrow \omega\chi_{b0}$ | 10.701           | $0.0^{+1.1}_{-0.0}$    | $< 16.6$                    |
| $e^+e^- \rightarrow \omega\chi_{b1}$ |                  | $0.0^{+2.1}_{-0.0}$    | $< 1.2$                     |
| $e^+e^- \rightarrow \omega\chi_{b2}$ |                  | $0.1^{+2.2}_{-0.1}$    | $< 2.5$                     |
| $e^+e^- \rightarrow \omega\chi_{b0}$ | 10.745           | $3.0^{+5.5}_{-4.7}$    | $< 11.3$                    |
| $e^+e^- \rightarrow \omega\chi_{b1}$ |                  | $68.9^{+13.7}_{-13.5}$ | $3.6^{+0.7}_{-0.7} \pm 0.5$ |
| $e^+e^- \rightarrow \omega\chi_{b2}$ |                  | $27.6^{+11.6}_{-10.0}$ | $2.8^{+1.2}_{-1.0} \pm 0.4$ |
| $e^+e^- \rightarrow \omega\chi_{b0}$ | 10.805           | $3.6^{+3.8}_{-3.1}$    | $< 11.4$                    |
| $e^+e^- \rightarrow \omega\chi_{b1}$ |                  | $15.0^{+6.8}_{-6.2}$   | $< 1.7$                     |
| $e^+e^- \rightarrow \omega\chi_{b2}$ |                  | $3.3^{+5.3}_{-3.8}$    | $< 1.6$                     |


Note:

$$\sigma_B(\chi_{b1}) = (0.76 \pm 0.16) \text{ pb}$$

$$\sigma_B(\chi_{b2}) = (0.29 \pm 0.14) \text{ pb}$$


at  $\sqrt{s} = 10.867$  (Belle)

arXiv:1408.0504v2 [hep-ex] 11 Sep 2014



# Observation of $Y(10753) \rightarrow \omega\chi_{bJ}$

- $\sigma_B$  fit features:
  - Model: PHSP + BW\*
  - $M$  and  $\Gamma$  of  $Y(10753)$  are fixed to 10752.7 MeV and 35.5 MeV.
- $\sigma_B$  fit result:
  - $\sigma_B$  enhancement near 10.753 GeV
  - No peak at  $\Upsilon(5S)$



- $\chi_{b1}/\chi_{b2}$  1: agrees with HQET.

arXiv:hep-ph/9908366v1 16 Aug 1999


$$* BW(\sqrt{s}) = \frac{\sqrt{12\pi}\Gamma_{ee}\mathcal{B}_f\bar{\Gamma}}{s - M^2 - iM\Gamma} \sqrt{\frac{\Phi_1(\sqrt{s})}{\Phi_2(M)}}$$

| $\Gamma_{ee}\mathcal{B}_f$  | Solution I<br>(constructive interference) | Solution II<br>(destructive interference) |
|---|---|---|
| $\Gamma_{ee}\mathcal{B}_f(\Upsilon(10753) \rightarrow \omega\chi_{b1})$ | $(2.01 \pm 0.38 \pm 0.46)$ eV             | $(0.63 \pm 0.39 \pm 0.20)$ eV             |
| $\Gamma_{ee}\mathcal{B}_f(\Upsilon(10753) \rightarrow \omega\chi_{b2})$ | $(0.53 \pm 0.46 \pm 0.15)$ eV             | $(1.32 \pm 0.44 \pm 0.55)$ eV             |

# Search for $X_b \rightarrow \omega\Upsilon(1S)$

- Search for resonances in  $\omega\Upsilon(1S)$  in each energy scan.
- $\omega\chi_{bJ}$  reflection is accounted.
- No evidence for  $X_b$  signal.
- \*  $\sigma_{X_b}^{UL}$  upper limit is set\*\* for each  $E_{CM}$  and  $M(X_b) \in [10.45, 10.65]$  GeV.

| $\sqrt{s}$ (GeV) | $M_{X_b}$ (GeV) | $\sigma_{X_b}^{UL}$ |
|------------------|-----------------|---------------------|
| 10.653           | 10.59           | 0.55                |
| 10.701           | 10.45           | 0.84                |
| 10.745           | 10.45           | 0.14                |
| 10.805           | 10.53           | 0.47                |



$$* \sigma_{X_b}^{UL} = \sigma_B^{UL}(e^+e^- \rightarrow \gamma X_b) \mathcal{B}(X_b \rightarrow \omega\Upsilon(1S))$$

\*\* at 90% Bayesian credibility

## Summary

---

# Charmonium(-like) studies at Belle II: from legacy prospects

- B-factories founded the XYZ studies, they are now complemented by other studies:
  - Many statistics-dominated B-decay modes covered by LHCb;
  - BES III energy scans extending range above 4.9 GeV;
- Still well-known for this legacy ( $X(3872)$  is still the most cited paper) and essential for full understanding of new states;
- Key future contributions:
  - Models with neutrals (e.g. neutral Z partners,  $\pi^0$  transitions/decays)
  - Unique double-charmonium ( $e^+e^- \rightarrow c\bar{c}c\bar{c}$ ) and two-photon ( $e^+e^- \rightarrow e^+e^-c\bar{c}$ ) production;
  - Statistics-dominated: results will come with the raise of integrated luminosity.

# The future of Bottomonium at Belle II

- Open questions Belle II can give answers to:
  - $\alpha_s$ -suppressed  $\Upsilon(nS)$  radiative transitions? (possible at Belle II only);
  - $Y_b$  nature;
  - $Z_b$  decomposition (only seen in  $\Upsilon(5S)$  decays so far). Other molecular states?
  - $\Upsilon(5S)$  mystery;
- Long-term non- $\Upsilon(4S)$  possibilities:
  - Revisit  $\Upsilon(6S)$  with  $10\times+$  statistics;
  - LFV/spectroscopy in  $\Upsilon(2S, 3S)$  decays;
  - Higher statistics scan of entire region and  $\Upsilon(5S)$ ;
  - $E_{CM}$  to  $\Lambda_b\bar{\Lambda}_b$  (requires further SuperKEKB upgrades).

## Golden Modes

$$e^+e^- \rightarrow \pi^+\pi^-\Upsilon(pS)(\rightarrow \ell^+\ell^-)$$

$B\bar{B}$  decomposition

$\pi^+\pi^-$  Dalitz

$$Y_b \rightarrow \omega\eta_b(1S)$$

$$Y_b \rightarrow \omega\chi_{bJ}(1P)$$

## Silver Modes

$$Y_b \rightarrow \pi^+\pi^-X \text{ (inclusive)}$$

$$Y_b \rightarrow \eta X \text{ (inclusive)}$$

$$Y_b \rightarrow \eta\Upsilon(1S, 2S)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow \eta'\Upsilon(1S)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow \Upsilon(1S) \text{ (inclusive)}$$

## Bronze Modes

$$Y_b \rightarrow \gamma X_b$$



$$Y_b \rightarrow \pi^0\pi^0\Upsilon(pS)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow KK(\phi)\Upsilon(pS)(\rightarrow \ell^+\ell^-)$$

$$Y_b \rightarrow \pi^0\pi^0X \text{ (inclusive)}$$

$$Y_b \rightarrow \pi^0X \text{ (incl. or excl.)}$$

...



Belle Phys. Rev. Lett. 100, 112001 (2008)



# Summary: Belle II take-off

- Huge experiment: 1100+ members, 123 institutions, 26 countries;
- Run time is scheduled until 2031+;
- First major upgrade is ongoing;
- Next long-term data collection period is to be launched in the next year.

