



L1 trigger efficiencies for $e^+e^- \rightarrow \tau^+\tau^-$ events

The Belle II Collaboration

Abstract

The Level 1 trigger efficiency has been studied in τ -pair events. The $ee \rightarrow \tau\tau \rightarrow 1 \times 3$ prong and 1×1 prong decay signatures are considered. These measurements were performed using the e^+e^- collision data recorded during the 2019a, 2019b and 2019c data taking periods. This document contains the approved plots and captions.

In this note we present the measurement of the Level 1 (L1) trigger efficiency for $e^+e^- \rightarrow \tau^+\tau^-$ events in 2019 data (periods 2019a, 2019b and 2019c). This data sample corresponds to a total integrated luminosity of 8.8 fb^{-1} .

The cross section for the process $e^+e^- \rightarrow \tau^+\tau^-$ is of the same order as the production of B meson pairs at the $\Upsilon(4S)$ resonance energy. In addition, τ lepton decays result in a wide variety of low multiplicity signatures involving tracks (e^\pm , μ^\pm or π^\pm) and ECL clusters (e^\pm , μ^\pm , π^\pm or π^0). Considering this, τ -pair events provide a good test bed for the performance of both the CDC and ECL triggers.

Three τ -pair decay modes are considered:

- *SM 1×1* — both τ leptons undergo Standard Model (SM) decay into one charged particle (1-prong). This can occur via the leptonic decay mode ($\tau^\pm \rightarrow \ell^\pm \nu_\ell \bar{\nu}_\tau$, $\ell=e, \mu$) or hadronic mode with one charged pion ($\tau^\pm \rightarrow \pi^\pm \nu_\tau + n\pi^0$).
- *SM 1×3* — one τ lepton undergoes SM 1-prong decay while the other undergoes SM decay into three charged pions (3-prong). The latter occurs through the hadronic decay mode with three charged pions ($\tau^\pm \rightarrow 3\pi^\pm \nu_\tau + n\pi^0$).
- *LFV 1×3* — one τ lepton decays undergoes SM 1-prong decay while the other undergoes Lepton Flavour Violating (LFV) decay into three muons. This targets one of the golden channels for LFV searches in τ decays at *Belle II* ($\tau \rightarrow 3\mu$). The strategy is to select SM 1×3 prong events that mimic the kinematics of the LFV process.

Six channels are considered according to whether the tracks are identified as electrons, muons or charged pions. For 1×1 prong decay we consider the $e\mu$, $\mu\pi$ and $\mu\mu$ channels. For 1×3 prong decay we have the $e - 3\pi$, $\mu - 3\pi$ and $\pi - 3\pi$ channels.

The following triggers have been studied:


- *CDC full tracks* — ffo and fff.
- *CDC short tracks* — fso, sso, ffs, fss and sss.
- *ECL* — hie, c4 and eclmumu.
- *ECL low multiplicity* — lml0, lml1, lml2, lml4, lml6, lml7, lml8, lml9, lml10 and lml12.

We define the efficiency of an individual CDC trigger as:


$$\epsilon = \frac{(\text{hie or c4 or eclmumu or lmlX}) \text{ and CDC trigger}}{\text{hie or c4 or eclmumu or lmlX}}, \quad (1)$$

and for an individual ECL trigger:

$$\epsilon = \frac{(\text{fff or ffo}) \text{ and ECL trigger}}{\text{fff or ffo}}, \quad (2)$$




(a)




(b)

FIG. 1: The overall L1 trigger efficiencies for SM $ee \rightarrow \tau\tau \rightarrow 1 \times 3$ prong events in the (a) combined and (b) individual channels. The data comes from the 2019a, 2019b and 2019c periods. The following trigger combinations are considered: ≥ 2 full tracks (ffo), ≥ 3 full tracks (fff), short tracks (fso or sso or ffs or fss or sss), ECL total energy (hie), ≥ 4 clusters (c_4), low multiplicity ≥ 3 clusters ($lml0$ or $lml12$), low multiplicity back-to-back clusters ($lml8$ or $lml9$ or $lml10$), low multiplicity high energy cluster ($lml1$ or $lml2$ or $lml4$ or $lml6$ or $lml7$) and ECL $\mu\mu$ ($eclmumu$). Statistical uncertainties are shown, although they are too small to be visible.




(a)




(b)

FIG. 2: The overall L1 trigger efficiencies for SM $ee \rightarrow \tau\tau \rightarrow 1 \times 1$ prong events in the (a) combined and (b) individual channels. The data comes from the 2019a, 2019b and 2019c periods. The following trigger combinations are considered: ≥ 2 full tracks (ffo), ≥ 3 full tracks (fff), short tracks (fso or sso or ffs or fss or sss), ECL total energy (hie), ≥ 4 clusters (c_4), low multiplicity ≥ 3 clusters ($lml0$ or $lml12$), low multiplicity back-to-back clusters ($lml8$ or $lml9$ or $lml10$), low multiplicity high energy cluster ($lml1$ or $lml2$ or $lml4$ or $lml6$ or $lml7$) and ECL $\mu\mu$ ($eclmumu$). Statistical uncertainties are shown, although they are too small to be visible.




(a)




(b)

FIG. 3: The efficiency of the ffo trigger for SM $ee \rightarrow \tau\tau \rightarrow 1 \times 1$ prong events. This trigger requires at least two full tracks, a track pair with $\Delta\phi > 90^\circ$ and an ECL Bhabha veto at L1. The efficiency is shown as a function of the (a) θ and (b) p_T of the track with minimum p_T . The data sample was collected during the 2019a, 2019b and 2019c periods. Statistical uncertainties are shown.




(a)




(b)

FIG. 4: The efficiency of the fff trigger for SM $ee \rightarrow \tau\tau \rightarrow 1 \times 3$ prong events. This trigger requires at least three full tracks at L1. The efficiency is shown as a function of the (a) θ and (b) p_T of the track with minimum p_T . The data sample was collected during the 2019a, 2019b and 2019c periods. Statistical uncertainties are shown.




(a)




(b)

FIG. 5: The gain in efficiency when using the ffo trigger in conjunction with short track triggers (ffo or fso or sso) for SM $ee \rightarrow \tau\tau \rightarrow 1 \times 1$ prong events. The efficiency is shown as a function of the (a) θ and (b) p_T of the track with minimum p_T . The data sample was collected during the 2019c period. Statistical uncertainties are shown.




(a)




(b)

FIG. 6: The gain in efficiency when using the fff trigger in conjunction with short track triggers (fff or ffs or fss or sss) for SM $ee \rightarrow \tau\tau \rightarrow 1 \times 3$ prong events. The efficiency is shown as a function of the (a) θ and (b) p_T of the track with minimum p_T . The data sample was collected during the 2019c period. Statistical uncertainties are shown.



(a)



(b)

FIG. 7: The efficiency of the *hie* trigger as a function of the ECL total cluster energy for (a) SM $ee \rightarrow \tau\tau \rightarrow 1 \times 1$ prong and (b) 1×3 prong events. This trigger has a 1 GeV total energy threshold and Bhabha veto requirement at L1. The data sample was collected during the 2019a, 2019b and 2019c periods. Statistical uncertainties are shown.







FIG. 8: The efficiency of the low multiplicity ≥ 3 cluster triggers ($lml0$ or $lml12$) for τ LFV-like events as a function of the signal-side ΔE . SM 1×3 prong events are considered that mimic the kinematics of the $\tau \rightarrow 3\mu$ LFV process. The data sample was collected during the 2019a, 2019b and 2019c periods. Statistical uncertainties are shown.



(a)



(b)



(c)

FIG. 9: The efficiency of the low multiplicity ≥ 3 cluster triggers ($lml0$ or $lml12$) for τ LFV-like events as a function of the signal-side ΔE . SM 1×3 prong events are considered that mimic the kinematics of the $\tau \rightarrow 3\mu$ LFV process. The tag-side track is identified as an (a) electron, (b) muon or (c) charged pion. The data sample was collected during the 2019a, 2019b and 2019c periods. Statistical uncertainties are shown.