



BELLE2-NOTE-PL-2019-ZZZ

Version 1.0

August 2, 2019

Approved plots for $J/\psi \rightarrow \ell^+\ell^-$ in Proc9

D. Ferlewicz,^{*} M. Milesi,[†] and P. Urquijo[‡]

School of Physics, University of Melbourne, Victoria, Australia

Abstract

Approved plots for the dilepton yields of $J/\psi \rightarrow \ell^+\ell^-$ for the analysis documented in BELLE2-NOTE-PH-2019-050.

^{*}Electronic address: daniel.ferlewicz@unimelb.edu.au

[†]Electronic address: marco.milesi@unimelb.edu.au

[‡]Electronic address: purquijo@unimelb.edu.au




FIG. 1: The dielectron invariant mass of $J/\psi \rightarrow e^-e^-$ candidates for an integrated luminosity of 2.63 fb^{-1} using the `basf2` software release `release-03-02-02` on the `hlt_hadron` skim. This data set includes Phase 3 physics runs only, excluding runs 916 – 1005 and 1216 – 1371 from experiment 7. The selection criteria are as follows: $|dr| < 2.0 \text{ cm}$, $|dz| < 5.0 \text{ cm}$, $p_{\text{lab}} > 0.1 \text{ GeV}/c$ and `electronID` > 0.95 for each electron candidate. A vertex fit using `TreeFitter` was applied, selecting candidates with a p-value > 0.001 . A bremsstrahlung correction was applied by adding the momentum and cluster energy of a photon with $E < 1.0 \text{ GeV}$ within a 5° cone of the electron candidate.

A Crystal Ball function summed with a Bifurcated Gaussian is used to model the signal and a first order polynomial is used to model the background.




FIG. 2: The dimuon invariant mass of $J/\psi \rightarrow \mu^+\mu^-$ candidates for an integrated luminosity of 2.63 fb^{-1} using the same environment and track selection as above, but with $\text{muonID} > 0.95$ for each muon candidate.

A Gaussian function summed with a Bifurcated Gaussian is used to model the signal and a first order polynomial is used to model the background.